CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 94, 2001

A
Adamczyk, J. J., 1445
Adamczyk, J. J., Jr., 284, 752, 1589
Adams, D. C., 780
Adams, L. C., 284, 1589
Adams, R. G., Jr., 455
Aggarwal, K. K., 979
Ahn, Y.-J., 157
Aihara-Sasaki, M., 785
Al-Deeb, M. A., 1353
Alderman, S. C., 1471
Aliniazee, M. T., 68
Altosaar, I., 271
Anderson, R. G., 430
Antilla, L., 315, 1012
Appel, A. G., 628
Argauer, R., 339
Armstrong, J. W., 302, 817
Arnaud, L., 552
Arthur, F. H., 772
Arzaluz, I. O., 1405
Ashfaq, M., 1053
Ashley, R. A., 455
Athanas, M. M., 892

B
Bacheler, J. S., 76
Bailey, W. D., 76
Baird, J. M., 1485
Bajwa, W. I., 68
Balbyshev, N. F., 1260
Barcenas, N., 958
Bartels, D. W., 197
Bartlett, A. C., 248
Baxendale, F., 743
Beers, E. H., 33, 1075
Behle, R., 1142
Behle, R. W., 1037, 1428
Bélanger, A., 167
Belliure, B., 439
Bengtsson, M., 911
Berisford, C. W., 112, 1122
Berkebile, D. R., 780
Berry, R. E., 1471
Bethke, J. A., 1031
Bierman, P., 1594
Bin, F., 648
Birdsall, S. L., 47
Bläske, V.-U., 1200
Blua, M. J., 1031, 1506
Boissot, N., 719
Bolin, P. C., 197
Bollich, P. K., 963
Bonning, B. C., 1045
Borgemeister, C., 1434
Bostanian, N., 167
Boucher, T. J., 455
Boughton, A. J., 1045
Bourguet, D., 1564
Boxler, D. J., 780
Bradley, J. R., Jr., 634
Braman, S. K., 129, 1187
Brandhorst-Hubbard, J. L., 628, 1452
Breisch, N. L., 492
Brenner, R. J., 1217
Brickle, D. S., 86
Brooks, H. L., 547
Brownie, C., 76
Brunner, J. F., 33, 1075, 1500
Buczkowski, G., 680, 1229
Buntin, G. D., 129, 1187
Burkness, E. C., 197
Burnell, A. M., 989
Buschman, L. L., 381, 929
Busey, P., 915
Butler, L., 761, 1128
Butler, M. D., 1471
Butrón, A., 277
Büyükgüzel, K., 21
Byrne, F. J., 524

C
Cabana, J., 1135
Cabras, P., 806
Calderone, N. W., 1022
Calkins, C. O., 1511
Cameron, P. J., 1104
Campbell, J. B., 780
Campora, C. E., 1193
Cao, J., 1547
Caprio, M. A., 698
Carbonell, E. A., 439
Cardona, C., 958
Cardwell, K. F., 1434
Carey, J. R., 971
Carlson, D. A., 1610
Carlson, J. R., 1584
Carrière, Y., 248, 315, 935, 1012, 1237, 1571
Cartea, M. E., 277
Carter, N. J., 476
Casaña-Giner, V., 898
Casey, D. G., 989
Casper, H., 1260
Castle, S. J., 1538
Celedonio, H., 1419
Chapin, J. W., 410
Chaufaux, J., 1564
Cheng, X., 14, 541
Cheng, X.-Y., 271
Chiasson, H., 167
Chirico, J., 1617
Chu, C.-C., 47
Coll, M., 27
Collins, H. L., 240, 1547
Conti, E., 648
Cooper, R. L., 1594
Coracini, M. D. A., 911
Corley, J. C., 811, 850
Costa, H. S., 1506
Coudron, T. A., 1344
Cowan, D., 1151
Crane, D. E., 116
Crisosto, C. H., 984
Crocker, R. L., 1452
Cui, H.-R., 271
Currie, R. W., 793

D
Daane, K. M., 291
d'Adamo, P., 850
Dalusky, M. J., 112
Davey, R. B., 1622
Davis, D. W., 197
Davis, F. M., 752
Davis, P. M., 586
Davis, R., 780
Dennehy, T. J., 315, 935, 1237, 1571
Ding, W., 1296
Dively, G. P., 1162
Doerr, M. D., 1075, 1500
Dosa, E. L., 1477
Dowd, P. F., 1067
Drummond, F. A., 609
Dunley, J. E., 33, 264, 1075

E
Earle, E. D., 240, 942, 1547
Edwards, J. K., 915
Eiras, A. E., 911
Ellers-Kirk, C., 315, 935, 1012, 1571
Elzen, G. W., 55, 357
English-Loeb, G., 1477
Erbilgin, N., 1113
Evans, T. A., 1602

F
Faber, B. A., 1085, 1090
Ferrari, J. A., 920
Ferster, B., 215
Fettig, C. J., 1122
Flanders, K. L., 628, 1452
Flinn, M., 558
Floris, I., 806

CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 94, 2001

A
Adamczyk, J. J., 1445
Adamczyk, J. J., Jr., 284, 752, 1589
Adams, D. C., 780
Adams, L. C., 284, 1589
Adams, R. G., Jr., 455
Aggarwal, K. K., 979
Ahn, Y.-J., 157
Aihara-Sasaki, M., 785
Al-Deeb, M. A., 1353
Alderman, S. C., 1471
Aliniazee, M. T., 68
Altosaar, I., 271
Anderson, R. G., 430
Antilla, L., 315, 1012
Appel, A. G., 628
Argauer, R., 339
Armstrong, J. W., 302, 817
Arnaud, L., 552
Arthur, F. H., 772
Arzaluz, I. O., 1405
Ashfaq, M., 1053
Ashley, R. A., 455
Athanas, M. M., 892

B
Bacheler, J. S., 76
Bailey, W. D., 76
Baird, J. M., 1485
Bajwa, W. I., 68
Balbyshev, N. F., 1260
Barcenas, N., 958
Bartels, D. W., 197
Bartlett, A. C., 248
Baxendale, F., 743
Beers, E. H., 33, 1075
Behle, R., 1142
Behle, R. W., 1037, 1428
Bélanger, A., 167
Belliure, B., 439
Bengtsson, M., 911
Berisford, C. W., 112, 1122
Berkebile, D. R., 780
Berry, R. E., 1471
Bethke, J. A., 1031
Bierman, P., 1594
Bin, F., 648
Birdsall, S. L., 47
Bläske, V.-U., 1200
Blua, M. J., 1031, 1506
Boissot, N., 719
Bolin, P. C., 197
Bollich, P. K., 963
Bonning, B. C., 1045
Borgemeister, C., 1434
Bostanian, N., 167
Boucher, T. J., 455
Boughton, A. J., 1045
Bourguet, D., 1564
Boxler, D. J., 780
Bradley, J. R., Jr., 634
Braman, S. K., 129, 1187
Brandhorst-Hubbard, J. L., 628, 1452
Breisch, N. L., 492
Brenner, R. J., 1217
Brickle, D. S., 86
Brooks, H. L., 547
Brownie, C., 76
Brunner, J. F., 33, 1075, 1500
Buczkowski, G., 680, 1229
Buntin, G. D., 129, 1187
Burkness, E. C., 197
Burnell, A. M., 989
Buschman, L. L., 381, 929
Busey, P., 915
Butler, L., 761, 1128
Butler, M. D., 1471
Butrón, A., 277
Büyükgüzel, K., 21
Byrne, F. J., 524

C
Cabana, J., 1135
Cabras, P., 806
Calderone, N. W., 1022
Calkins, C. O., 1511
Cameron, P. J., 1104
Campbell, J. B., 780
Campora, C. E., 1193
Cao, J., 1547
Caprio, M. A., 698
Carbonell, E. A., 439
Cardona, C., 958
Cardwell, K. F., 1434
Carey, J. R., 971
Carlson, D. A., 1610
Carlson, J. R., 1584
Carrière, Y., 248, 315, 935, 1012, 1237, 1571
Cartea, M. E., 277
Carter, N. J., 476
Casaña-Giner, V., 898
Casey, D. G., 989
Casper, H., 1260
Castle, S. J., 1538
Celedonio, H., 1419
Chapin, J. W., 410
Chaufaux, J., 1564
Cheng, X., 14, 541
Cheng, X.-Y., 271
Chiasson, H., 167
Chirico, J., 1617
Chu, C.-C., 47
Coll, M., 27
Collins, H. L., 240, 1547
Conti, E., 648
Cooper, R. L., 1594
Coracini, M. D. A., 911
Corley, J. C., 811, 850
Costa, H. S., 1506
Coudron, T. A., 1344
Cowan, D., 1151
Crane, D. E., 116
Crisosto, C. H., 984
Crocker, R. L., 1452
Cui, H.-R., 271
Currie, R. W., 793

D
Daane, K. M., 291
d'Adamo, P., 850
Dalusky, M. J., 112
Davey, R. B., 1622
Davis, D. W., 197
Davis, F. M., 752
Davis, P. M., 586
Davis, R., 780
Dennehy, T. J., 315, 935, 1237, 1571
Ding, W., 1296
Dively, G. P., 1162
Doerr, M. D., 1075, 1500
Dosa, E. L., 1477
Dowd, P. F., 1067
Drummond, F. A., 609
Dunley, J. E., 33, 264, 1075

E
Earle, E. D., 240, 942, 1547
Edwards, J. K., 915
Eiras, A. E., 911
Ellers-Kirk, C., 315, 935, 1012, 1571
Elzen, G. W., 55, 357
English-Loeb, G., 1477
Erbilgin, N., 1113
Evans, T. A., 1602

F
Faber, B. A., 1085, 1090
Ferrari, J. A., 920
Ferster, B., 215
Fettig, C. J., 1122
Flanders, K. L., 628, 1452
Flinn, M., 558
Floris, I., 806

Follett, P. A., 1005, 1292
Francke, W., 911
Franqui, R. A., 853
Freidberg, A., 1059
Frelichowski, J. E., Jr., 1249
Furlong, M. J., 344

G

Gahlhoff, J. E., Jr., 486
Gandía-Balaguer, A., 898
Ganz, P. J., 1584
Gao, M.-W., 271
Garau, V. L., 806
Garcia-Marí, F., 1558
Gardner, J., 1485
Garrido-Vivas, A., 898
Garza, R., 958
Gates, R. S., 430
Gavrilovic, V., 1378
Geden, C. J., 1610
Geiger, C. A., 291
George, J. E., 1622
Georghiou, G. P., 920
Gibb, T. J., 145
Gill, B., 558
Giroux, P. Y., 1170
Gore, J., 1445
Gould, F., 76
Grace, J. K., 1193
Granett, J., 138
Gray, M. E., 98, 529
Gray, S. M., 410
Greene, J. K., 403
Groden, E., 344
Guerrero, F. D., 714
Guertal, E. A., 1452
Guertin, C., 1135
Guillaume, R., 719
Guo, B. Z., 564
Guse, C. A., 529

H

Haack, R. A., 422
Halbert, S. E., 410
Hall, D. B., 666
Haller, S., 315, 1012
Hallett, R. H., 1177
Halliday, W. R., 397
Hammond, A. M., 1285
Hammond, P. C., 1471
Hammond, R. B., 1594
Hanley, A. M., 586
Harbo, J. R., 1319
Hardee, D. D., 284, 1589
Harris, J. W., 1319
Harris, M. K., 180
Haubruge, E., 552
He, F., 929
Heath, R. R., 971
Heaton, G. C., 422
Held, D. W., 60, 430
Hellmich, R. L., 1369
Henderson, G., 1532
Hendrichs, J., 971
Heng-Moss, T., 743
Henneberry, T. J., 47, 1538
Herbert, D. A., Jr., 634
Herman, T. J. B., 1104
Hermoso de Mendoza, A., 439
Hernández-Alamós, M. M., 898
Hertel, H., 1200
Hewa-Kapuge, S., 826
Higbee, B. S., 1511
Higgins, R. A., 929
Higley, L. G., 1369
Hines, R. L., 190
Hinton, J. L., 1308
Hoffmann, A. A., 826
Hoffmann, M. P., 1485
Horowitz, A. R., 1059
Hower, A. A., 950
Hu, C., 271
Huang, X., 860
Hung, R. W., 123
Hunt, D. W. A., 204
Hunt, T. E., 1369
Hutchison, W. D., 190, 197

I

Ihrig, R. A., 634
Inglis, G. D., 362
Ingram, D., 388
Ishaaya, I., 1059

J

Jackson, C. G., 817
Jackson, J., 445
Jadhav, D., 253
James, R. R., 357
Jansson, R. K., 264
Jones, R. W., 1405
Jones, V. P., 785
Jordan, R. A., 123
Jordan, T. M., 123
Juvik, J. A., 1249
Jyoti, J. L., 942

K

Kahn, A. D., 1500
Kairu, E. W., 449
Kakouli-Duarte, T., 989
Kamble, S. T., 499
Kammlah, D. M., 714
Kang, J.-G., 666
Katsoyannos, B. I., 971
Kaufman, P. E., 593, 998
Kawamura, S., 1340
Kennedy, G. G., 76, 1162, 1268
Kevan, P. G., 172, 462
Khan, Z. R., 449
Kim, Y.-J., 157
Kindler, S. D., 547
Klein, M. G., 1151
Klepzig, K. D., 1113
Klingeman, W. E., 129
Klingeman, W. E., III, 1187
Klucas, R., 743
Knight, A. L., 264
Knighten, K. S., 752
Knowles, C. O., 868, 874, 884
Koehler, P. G., 486, 1209
Komaravalli, S. R., 586
Kono, Y., 223
Kopanic, R. J., Jr., 686, 1229
Korie, S., 1434
Kotze, A. C., 1243
Kouassi, K. C., 1135
Kouloussis, N. A., 971
Kovach, J., 1477
Kovaleski, A., 911
Kranthi, K. R., 253
Kranthi, S., 253
Krecek, J., 915
Krivenko, A. J., Jr., 123
Krupke, C. H., 1500
Kuenen, L. P. S., 1022
Kuhar, T. P., 1485
Kumar, S., 979
Kunkel, B. A., 60

L

LaBonte, D. R., 1285
Ladonni, H., 694
Lafta, A. M., 1260
Laine, R. A., 1532
Lam, W.-K. F., 1459
Lamp, W. O., 93
LaPointe, S. L., 853
Laverty, T. M., 172, 462
Lawrence, A. M., 362
Lawrence, R. K., 422
Lax, A. R., 1217
Layton, M. B., 388
Lee, S.-E., 706
Lees, E. M., 706
Leonard, B. R., 1445
Levine, E., 529, 1594
Lewis, L. C., 617, 1045
Li, L.-S., 1296
Li, R. G., 564
Li, Y.-X., 1547
Liang, D., 686
Liburd, O. E., 905, 1142
Linduska, J., 1162
Linscombe, S. D., 963
Liu, Y.-B., 248, 315, 935, 1237, 1571
Löfqvist, J., 911
Lonergan, G., 658
Long, S. J., 593, 998
Lord, J. C., 367
Lorenzen, J. H., 1260
Lorenzetti, F., 1135
Lozada, M., 850
Lynch, R. E., 564

M
Machuka, J., 724
Mack, T. P., 860
Maharajh, B., 915
Maily, W., 388
Maistrello, L., 1532
Maklakov, A., 1059
Malvar, R. A., 277
Mangan, R. L., 150, 1419
Mankin, R. W., 853, 1452
Mannion, C. M., 1151
Mao, L., 1285
Margolies, D. C., 381
Markwell, J., 743
Martin, P. A. W., 892
Mastro, V. C., 116
Mauffette, Y., 1135
May, O. L., 403
Mbata, G. N., 1302
McCravy, K. W., 112, 1122
McGuire, M. R., 1037, 1142
McInnis, D. O., 1413
McLane, W., 1151
McNew, R. W., 1053
McQuate, G. T., 39, 817, 1005
Meinke, L. J., 106
Meissner, H. E., 1526
Mengod-Puerta, C., 898
Metz, T. D., 240
Meyer, S., 935
Meyer, S. K., 248, 1237
Miller, F., 162, 445
Miller, G. T., 984
Miller, J. A., 1622
Miller, J. R., 1170
Minelli, E. V., 806
Mitchell, P. D., 599
Miyo, T., 223
Montoya, P., 1419
Moon, R. D., 1308
Morales-Ramos, J. A., 506, 516
Morandin, L. A., 172, 462
Moreno, D. S., 150, 1419
Morgan, D. J. W., 1506
Morris, T. F., 455
Mowry, T. M., 1332
Moyseenko, J., 1151
Musabyimana, T., 449
Myers, J. H., 831

N
Nalyanya, G., 208, 686
Nasr, M. E., 332
Natwick, E. T., 47
Nauen, R., 1577
Nault, B. A., 1162
Nava-Camberos, U., 180
Ndemah, R., 1434
Neilsen, K., 658
Nevo, E., 27
Ni, X., 743
Nielsen, G. R., 93
Nordenfors, H., 1617
Nowak, D. J., 116
Nowak, J. T., 1122

O
Obrycki, J. J., 617
Oehler, D. D., 1622
Oetting, R. D., 666
Ogol, C. P. K. O., 449
Oguma, Y., 223
Okeola, O. G., 724
Oliver, J. B., 1151
Omer, A. D., 138
O'Neal, M. E., 98
Onstad, D. W., 529
Ordás, A., 277
O'Rourke, S. D., 1308
Osbrink, W. L. A., 1217
Otis, G. W., 332
Ottea, J. A., 1464, 1553

P
Papadopoulos, N. T., 971
Park, N.-J., 499
Pasek, J. E., 116
Patin, A. L., 935, 1237, 1571
Pearsall, I. A., 831
Peck, S. L., 39, 817
Pedersen, B., 315, 1012
Pedigo, L. P., 1459
Pelletier, Y., 572
Peng, H., 658
Pernal, S. F., 793
Peterson, J. K., 1285
Petrice, T. R., 1128
Phillips, P. A., 1085, 1090
Phillips, T. W., 1302
Phoofolo, M. W., 617
Pierce, C. M. F., 1361
Pilcher, C. D., 730
Piñero, J. C., 1386
Plaisted, D., 564
Plaugher, C. D., 1128
Polavarapu, S., 658
Poliquin, A., 167
Potter, D. A., 60, 430
Pound, J. M., 1622
Prabhaker, N., 1538
Prajapati, V., 979
Pree, D. J., 476
Price, B. D., 586
Primo-Millo, J., 898
Primo-Yúfera, E., 898
Prokopy, R. J., 1142, 1386
Pruett, J. H., 714
Puche, H., 1398

Q
Quebedeaux, B., 93
Quisenberry, S. S., 743

R
Rabinovich, M., 811
Raffa, K. F., 844, 1113
Ranger, C. M., 950
Ratcliffe, S., 98
Reagan, T. E., 766, 1464, 1553
Real, V., 439
Reckziegel, A., 911
Redak, R. A., 1031, 1506
Reding, M. E., 33
Redmer, J. S., 844
Reed, J. P., 397
Reed, J. T., 640
Reeder, N. L., 1584
Reese, J. C., 558
Reissig, W. H., 468, 673
Reuter, G. S., 326
Rhainds, M., 1477
Rice, M. E., 730
Rice, W. C., 963
Richmond, D. S., 1183
Riedell, W. E., 599
Riley, D. G., 180
Ring, D. R., 766
Ripault, G., 623
Ritter, D., 47
Rivault, E., 623
Rodriguez, L. M., 1464, 1553
Roe, R. M., 76
Rojas, M. G., 506, 516
Roulston, T. H., 511
Roush, R. T., 240
Russell, D., 253
Rutz, D. A., 593, 998

S
Sackmann, P., 811
Sagredo, B., 1260
Sales, N., 1243
Salzman, R. A., 1280
Samson, P. R., 1097
Sanchez-Arroyo, H., 1209
Sanxter, S. S., 1292
Sarath, G., 743
Satta, A., 806
Saunders, C. W., 1584
Saxena, R. C., 449
Schal, C., 208, 680, 686, 1229
Scheffer, S. J., 1177
Scheffrahn, R. H., 215, 915
Scherer, P. N., 215
Schexnayder, H. P., Jr., 766
Schmidt, D. R., 1308
Schroder, R. F. W., 892
Schulthess, F., 1434
Schulze, J. J., 123
Schulze, T. L., 123
Schwartz, P., 1485
Scott-Dupree, C. D., 332
Sedlacek, J. D., 586
Seguin, M., 1564
Sequeira, R. A., 116

Shapiro, M., 1, 339
Shapiro-Ilan, D. I., 7
Shearer, P. W., 233
Shelly, T. E., 1413
Shelton, A. M., 14, 240, 541, 942, 1485, 1547
Sheppard, W. S., 1324
Shetlar, D. J., 1183
Shipp, J. L., 579
Shu, Q.-Y., 271
Shufran, R. A., 547, 929
Siegfried, B. D., 1564
Silverman, J., 511, 1526
Sims, M. A., 935, 1237, 1571
Singh, S. P., 958
Skoda, S. R., 780
Sloderbeck, P. E., 547, 929
Smith, C. M., 558
Smith, D. B., 640
Smith, D. M., 410
Smith, J. J., 1378
Smitley, D. R., 1378
Snook, M. E., 564
Solter, L. F., 1361
Sorenson, C. E., 76
Soto, A., 1558
Speese, J., III, 1162
Spencer, J. L., 529
Spivak, M., 326
Stark, J. D., 817
Starkey, S. R., 929
Steffey, K. L., 98
Stelinski, L. L., 905, 1142
Stewart, S. D., 388, 752
Storer, N. P., 1268
Story, R. N., 1285
Stout, M. J., 963
Stoyenoff, J. L., 1492
Strange, J. P., 1324
Strazanac, J., 761
Strazanac, J. S., 1128
Streett, D. A., 1037
Stubbs, C. S., 609
Stumpf, N., 1577
Su, N.-Y., 915, 1398, 1518
Sullivan, M. J., 86, 403
Sumerford, D. V., 284
Swanson, J. J., 1564
Szele, A., 1113

T

Tabashnik, B. E., 248, 315, 935, 1237, 1571
Tai, G. C. C., 572
Takada, Y., 1340
Takamori, H., 223
Tamez-Guerra, P., 1037
Tanaka, T., 1340
Tang, J. D., 240
Tasei, J. N., 623
Temple, C. A., 1511
Thomas, G. D., 780
Thomas, J. S., 410
Thoms, E. M., 215
Thorne, B. L., 492
Tian, X., 1260
Toscano, N. C., 524, 1538
Trimble, R. M., 476
Tripathi, A. K., 979
Troullier, J., 806
Tsai, J. H., 1296
Turnipseed, S. G., 86, 403

U

Urías-López, M. A., 106
Usmani, K. A., 233, 868, 874, 884

V

Vales, M. I., 277
Valles, S. M., 1209
Van Duyn, J. W., 634, 1268
van Iersel, M. W., 666
Vargas, R. I., 817
Vera, J., 958
Vernon, R. S., 204
Vilela, E. F., 911
Villanueva-Jiménez, J. A., 1004
Vincent, C., 167
Visser, D., 1177

W

Waldron, J. K., 593
Waldstein, D. E., 468, 673
Walker, G. P., 1104
Walker, K. R., 373
Walker, M. A., 138
Wallace, A. R., 1104
Wallin, K. F., 844
Wang, C., 761
Wang, J.-J., 1296
Wang, K., 579
Wang, Z., 93
Wanjari, R., 253
Ware, G., 162, 445
Warnock, D. F., 197
Weinzierl, R. A., 1361
Welter, S. C., 373
Whitworth, R. J., 929
Widstrom, N. W., 564
Wijesekara, A., 1177
Wilde, G. E., 547, 929, 1353
Williams, M. R., 388
Willott, E., 315
Wirth, M. C., 920
Witkowski, J. F., 1369
Wittmeyer, J. L., 1344
Witzgall, P., 911
Wright, S., 1142
Wright, S. E., 1386

X

Xia, Y.-W., 271
Xu, J., 14, 541

Y

Yang, X., 381
Yao, H.-W., 271
Yarom, I., 1059
Yawetz, A., 1059
Ye, G.-Y., 271
Yee, W. L., 1085, 1090
Yokoyama, V. Y., 984
Yoo, J.-K., 157
Young, L. J., 1369
Young, S. Y., 1053

Z

Zavala, J. L., 1419
Zenger, J. T., 145
Zhang, Z. J., 564
Zhao, J. Z., 240, 1547
Zhao, Z.-M., 1296
Zhu, K. Y., 381, 929, 1353
Zhu-Salzman, K., 1280

CUMULATIVE SUBJECT INDEX
Journal of Economic Entomology
Volume 94, 2001

1

1,8-cineole,
antifeedant, 979
growth inhibition, 979
toxicity, 979
Tribolium castaneum, 979

A

a1-sh2,
favonoid, 564
Helicoverpa zea, 564
marker-assisted selection, 564
Zea mays, 564
abamectin,
avocado, 1090
helicopter application, 1090
Scirtothrips perseae, 1090
spray volumes, 1090
Acalymma vittatum,
Citrullus lanatus, 892
Diabrotica undecimpunctata, 892
Hawkesbury watermelon, 892
southern corn rootworm, 892
xanthene dye, 892
Acarapis woodi,
Apis mellifera, 332
bee breeding, 332
honey bee, 332
mite resistance, 332
tracheal mite, 332
acaricide,
Artemisia absinthium, 167
essential oil extracts, 167
Tanacetum vulgare, 167
twospotted spider mite, 167
Acephate,
Bacillus thuringiensis variety *kurstaki*, 112
biological insecticides, 112
parasitism, 112
Rhyacionia frustrana, 112
tebufenozide, 112
acetylcholinesterase insensitivity,
chlorpyrifos, 524
insecticide resistance, 524
methomyl, 524
Spodoptera exigua, 524
AChE,
Blattella germanica, 499
chlorpyrifos, 499
esterase, 499
resistance, 499
AcMNPV,
AfMNPV, 1037
field activity, 1037
formulation, 1037
Lepidoptera, 1037
solar stability, 1037
acoustic sampling,
geostatistics, 1452
soil insects, 1452
Acrosternum hilare,
Euschistus servus, 403
Nezara viridula, 403
action threshold, 190, 1332
cabbage, 190
insecticidal control, 1332
integrated pest management, 190
Myzus persicae, 1332
potato leafroll virus, 1332
Trichoplusia ni, 190
vector, 1332
adult mortality,
corn rootworm, 599
Diabrotica barberi, 599
Diabrotica virgifera virgifera, 599
dispersal, 599
larval survival, 599
aesthetic threshold,
azalea lace bug, 1187
hybrid economic injury level, 1187
Rhododendron spp., 1187
Stephanitis pyrioides, 1187
AfMNPV,
AcMNPV, 1037
field activity, 1037
formulation, 1037
Lepidoptera, 1037
solar stability, 1037
African yam bean,
Clavigralla tomentosicollis, 724
insect bioassays, 724
lectins, 724
Sphenostylis stenocarpa, 724
aggregation pheromone, 1386, 1500
Conotrachelus nenuphar, 1386
Euschistus conspersus, 1500
fruit volatiles, 1386
Hemiptera, 1500
monitoring traps, 1386
orchard, 1500
Pentatomidae, 1500
plum curculio, 1386
trapping, 1500
Agromyzidae,
introduced species, 1177
invasive species, 1177
molecular identification, 1177
pea leafminer, 1177
polymerase chain reaction-restriction fragment-length polymorphism, 1177
Agrotis ipsilon, 1045, 1183
baculovirus, 1045
black cutworm, 1183
endophyte, 1183
M2R, 1045
optical brightener, 1045
perennial ryegrass, 1183
Aleurothrixus floccosus,
bioassay, 1558
insecticide resistance, 1558
alfalfa, 93, 950
Empoasca fabae, 93, 950
glandular trichomes, 950
host plant resistance, 950
Medicago sativa, 950
phloem translocation, 93
potato leafhopper, 950
alpha-ionol,
Bactrocera latifrons, 39
cade oil, 39
parapheromone, 39
Alphitobius diaperinus,
Dermestes maculatus, 1610
hide beetle, 1610
lesser mealworm, 1610
litter beetles, 1610
mechanical barrier, 1610
alternative treatment efficacy,
Bactrocera dorsalis, 1005
Cryptophlebia, 1005
pest risk management, 1005
quarantine treatment, 1005
amplified fragment-length polymorphism,
Ceratitis capitata, 989
Ceratitis rosa, 989
DNA probes, 989
silver staining, 989
tephritids, 989
Anastrepha ludens,
insecticide, 150
Mexican fruit fly, 150
photo active dye, 150
Angoumois grain moth,
Indian meal moth, 586
stored grain, 586
transgenic grain, 586
Anoplophora glabripennis,

urban forestry, 116
urban forests, 116
Anthonomus grandis,
ecology, 1405
herbivory, 1405
host plants, 1405
antibiosis, 558, 958
antixenosis, 558
Apion godmani, 958
goatgrass, 558
host plant resistance, 958
hypersensitivity, 958
Phaseolus vulgaris, 958
Schizaphis graminum, 558
tolerance, 558
wheat, 558
antifeedant,
1,8-cineole, 979
growth inhibition, 979
toxicity, 979
Tribolium castaneum, 979
antixenosis,
antibiosis, 558
goatgrass, 558
Schizaphis graminum, 558
tolerance, 558
wheat, 558
ants,
Argentine ant, 511
bait, 511
Ants,
bait traps, 761
forest, 761
Formicidae, 761
ants, 145, 511
gel, 511
halofenozide, 145
imidacloprid, 145
Japanese beetles, 145
Linepithema humile, 511
liquid, 511
Ants,
pitfall traps, 761
ants,
predators, 145
turfgrass, 145
aphid–plant interactions,
Aphis gossypii, 27
cotton, 27
fecundity, 27
morphology, 27
nitrogen fertilization, 27
Aphis gossypii, 27, 439, 1492
aphid–plant interactions, 27
Citrus clementina, 439
cotton, 27
economic injury level, 439
economic threshold, 439
fecundity, 27
greenhouse, 1492
hibiscus, 1492
horticulture, 1492
morphology, 27
nitrogen fertilization, 27
pest management, 1492
wash, 1492
apical damage,
introduced crop, 648
kenaf, 648
Lygus pratensis, 648
Lygus rugulipennis, 648
Apion godmani,
antibiosis, 958
host plant resistance, 958
hypersensitivity, 958
Phaseolus vulgaris, 958
Apis mellifera, 326, 332, 609, 793, 1022, 1324
Acarapis woodi, 332
bee breeding, 332
Bombus impatiens, 609
bumble bee, 609
flight room, 793
fluvalinate, 1324
foraging, 793
honey bee, 332, 609, 1022, 1324
host-parasite, 1022
hygienic behavior, 326
insect rearing, 793
insect vision, 793
lowbush blueberry, 609
mite resistance, 332
parasitic mite, 1022
pollination, 609
tracheal mite, 332
treatment threshold, 1324
Varroa destructor, 1022, 1324
Varroa destructor mite resistance, 326
Varroa jacobsoni, 1022, 1324
Apis mellifera,
breeding, 1319
resistance, 1319
selection, 1319
Varroa destructor, 1319
apple, 68, 468, 1511
Bacillus thuringiensis, 68
chemical insecticides, 68
Choristoneura rosaceana, 468
codling moth, 1511
Cydia pomonella, 1511
ecotoxicology, 68
exposure, 468
overwintering, 1511
pear, 1511
pest management, 1511
population dynamics, 68
residue, 468
spiders, 68
tebufenozide, 468
apples,
Choristoneura rosaceana, 673
insect growth regulator, 673
obliquebanded leafroller, 673
tebufenozide, 673
areawide control,
bait sprays, 817
biological control, 817
Ceratitis capitata, 817
Fopius arisanus, 817
areawide pest management,
carbaryl, 929
Diabrotica virgifera virgifera, 929
insecticide resistance, 929
Argentine ant,
ants, 511
bait, 511
gel, 511
Linepithema humile, 511
liquid, 511
Argyrotaenia citrana,
leafrollers, 373
mating disruption, 373
Pandemis pyrusana, 373
pheromones, 373
aromatic cedar mulch,
cultural control, 1526
landscaping, 1526
Linepithema humile, 1526
Tapinoma sessile, 1526
arsenic,
borate, 215
Incisitermes snyderi, 215
Spinosad, 215
Artemisia absinthium,
acaricide, 167
essential oil extracts, 167
Tanacetum vulgare, 167
twospotted spider mite, 167
arthropods,
bedding plants, 430
floriculture, 430
greenhouse, 430
hypoxia, 430
attractant,
carbon dioxide, 860
Elasmopalpus lignosellus, 860
attractants, 208, 686, 898
baits, 208, 686
Blattella germanica, 208, 686
Ceratitis capitata, 898
cockroach, 686
Mediterranean fruit fly, 898
Supella longipalpa, 686
traps, 208
automated termite monitoring system,
colony elimination, 1518
datalogger, 1518
Isoptera, 1518
aversion,
imidacloprid, 492
learning, 492
Premise, 492

repellency, 492
Reticulitermes virginicus, 492
avocado, 1085, 1090
abamectin, 1090
helicopter application, 1085, 1090
sabadilla, 1085
Scirtothrips perseae, 1085, 1090
spray volumes, 1085, 1090
Azadirachta indica,
banana, 449
Cosmopolites sordidus, 449
neem, 449
azalea lace bug, 129, 1187
aesthetic threshold, 1187
hybrid economic injury level, 1187
Rhododendron spp., 129, 1187
Stephanitis pyrioides, 129, 1187
azinphosmethyl,
bioassay, 233
Grapholita molesta, 233
resistance monitoring, 233
synergism, 233

B

B. thuringiensis cotton,
Bacillus thuringiensis, 1237
Cry1Ac, 1237
Pectinophora gossypiella, 1237
resistance, 1237
B.t. cotton,
Bollgard, 86
bollworm, 86
Helicoverpa zea, 86
insecticides, 86
Bacillus thuringiensis quantification,
dual-toxin, 1589
ELISA, 1589
transgenic cotton varieties, 1589
Bacillus sphaericus,
Bacillus thuringiensis, 920
Culex pipiens, 920
microbial control, 920
population variation, 920
Bacillus thuringiensis, 68, 76, 197, 240, 248, 271, 315, 397, 698, 730, 920, 935, 1012, 1053, 1067, 1135, 1237, 1268, 1361, 1547, 1564, 1571
apple, 68
B. thuringiensis cotton, 1237
Bacillus sphaericus, 920
bioassay, 1361
bollworm, 76
chemical insecticides, 68
Chilo suppressalis, 271
corn earworm, 1268
cry1Ab gene, 271
Cry1Ac, 248, 1237
Cry1C, 1547
Cry9C, 397
Culex pipiens, 920
diapause, 935, 1012
Diatrea grandiosella, 397
dispersal, 315, 698
ecotoxicology, 68
European corn borer, 397
fitness costs, 935
fumonisin, 1067
Fusarium, 1067
genetically engineered crops, 730
geographical information system, 315
heat units, 1012
Helicoverpa, 1067
Helicoverpa zea, 76, 197
Heliothis virescens, 76
host plant-mediated effect, 1135
inheritance, 248
insect pathogen, 1361
insecticide resistance, 76, 698
larval development, 1361
LC_{50}, 1135
Lepidoptera, 1564
local adaptation, 698
Malacosoma disstria, 1135
maternal effects, 1571
microbial control, 920
Nosema pyrausta, 1361
Ostrinia, 1067
Ostrinia nubilalis, 197, 397, 730, 1361, 1564
Pectinophora gossypiella, 248, 315, 935, 1012, 1237, 1571
Plutella xylostella, 240, 1547
population dynamics, 68
population variation, 920
Pseudoplusia includens, 1053
quaking aspen, 1135
resistance, 240, 1237, 1547, 1564
resistance costs, 1571
resistance evaluation, 271
resistance management, 248, 315, 1012
Scirpophaga incertulas, 271
southwestern corn borer, 397
soybean looper, 1053
spiders, 68
sublethal effects, 1268
sugar maple, 1135
tobacco budworm, 76
transgenic corn, 1564
transgenic cotton, 248, 315, 935, 1012, 1053, 1571
transgenic crops, 698, 1268
transgenic plants, 240, 1547
transgenic rice, 271
transgenic sweet corn, 197
Zea mays, 730
Bacillus thuringiensis quantification,
enzyme-linked immunosorbent assay, 284
transgenic varieties, 284
Bacillus thuringiensis variety *kurstaki*,
Acephate, 112
biological insecticides, 112
parasitism, 112
Rhyacionia frustrana, 112
tebufenozide, 112
Bactrocera dorsalis,
alternative treatment efficacy, 1005
Cryptophlebia, 1005
pest risk management, 1005
quarantine treatment, 1005
Bactrocera latifrons,
alpha-ionol, 39
cade oil, 39
parapheromone, 39
baculovirus,
Agrotis ipsilon, 1045
M2R, 1045
optical brightener, 1045
bait, 506, 511, 1229
ants, 511
Argentine ant, 511
Blattella germanica, 1229
chitin synthesis inhibitors, 506
control, 506
coprophagy, 1229
gel, 511
horizontal toxicant transfer, 1229
Linepithema humile, 511
liquid, 511
subterranean termite, 506
bait sprays,
areawide control, 817
biological control, 817
Ceratitis capitata, 817
Fopius arisanus, 817
bait traps,
ants, 761
forest, 761
Formicidae, 761
pitfall traps, 761
baits, 208, 686
attractants, 208, 686
Blattella germanica, 208, 686
cockroach, 686
Supella longipalpa, 686
traps, 208
banana, 302, 449
Azadirachta indica, 449
Cosmopolites sordidus, 449
Musa acuminata, 302

neem, 449
quarantine, 302
tephritid fruit flies, 302
trapping, 302
bark beetle,
exotic pest, 422
Scolytidae, 422
Tomicus piniperda, 422
barley yellow dwarf,
Rhopalosiphum padi, 410
Rhopalosiphum rufiabdominalis, 410
Schizaphis graminum, 410
Sitobion avenae, 410
wheat, 410
baseline susceptibility,
Bemisia argentifolii, 1538
resistance monitoring, 1538
whiteflies, 1538
bean leaf beetle,
Cerotoma trifurcata, 1459
glabrous, 1459
pubescence, 1459
trichome, 1459
Beauveria bassiana, 344, 357, 367
Bemisia argentifolii, 357
Coleoptera, 367
Colorado potato beetle, 344
cyromazine, 344
diatomaceous earth, 367
imidacloprid, 344, 357
insect pathogens, 357
microbial control, 357
mycopathogen, 367
starvation, 344
stored grain, 367
synergism, 344, 367
whiteflies, 357
bedding plants,
arthropods, 430
floriculture, 430
greenhouse, 430
hypoxia, 430
bee breeding,
Acarapis woodi, 332
Apis mellifera, 332
honey bee, 332
mite resistance, 332
tracheal mite, 332
beet armyworm,
host plant resistance, 1249
Lycopersicon hirsutum, 1249
sesquiterpene, 1249
tomato, 1249
tomato fruitworm, 1249
beetles,
exposure, 772
hydroprene, 772
morphological effects, 772
toxicity, 772
treated surfaces, 772
behavioral control,
deployment strategies, 905
imidacloprid-treated spheres, 905
Rhagoletis mendax, 905
Bemisia,
Bemisia argentifolii, 47
CC trap, 47
suction trap, 47
yellow sticky card trap, 47
Bemisia argentifolii, 47, 180, 357, 666, 1538
baseline susceptibility, 1538
Beauveria bassiana, 357
Bemisia, 47
cantaloupe, 180
CC trap, 47
economic injury levels, 180
Euphorbia pulcherrima, 666
imidacloprid, 357
insect pathogens, 357
insecticide efficacy, 666
irrigation method, 666
microbial control, 357
resistance monitoring, 1538
suction trap, 47
whiteflies, 357, 1538
yellow sticky card trap, 47
beneficial insect,
fecundity, 1344
intrinsic rate of increase, 1344
Podisus maculiventris, 1344
predator, 1344
rearing, 1344
bioassay, 55, 233, 1075, 1308, 1361, 1558
Aleurothrixus floccosus, 1558
azinphosmethyl, 233
Bacillus thuringiensis, 1361
biological control, 1075
compost, 1308
Geocoris punctipes, 55
Grapholita molesta, 233
house fly, 1308
insect pathogen, 1361
insecticide, 55
insecticide resistance, 1558
larval development, 1361
larval nutrition, 1308
Musca domestica, 1308
Nosema pyrausta, 1361
Orius insidiosus, 55
Ostrinia nubilalis, 1361
pesticides, 1075
poultry manure, 1308
resistance monitoring, 233
sublethal, 1075
synergism, 233
toxicity, 1075
biological control, 7, 593, 617, 817, 998, 1075, 1340
areawide control, 817
bait sprays, 817
bioassay, 1075
Ceratitis capitata, 817
Coleomegilla maculata, 617
Curculio caryae, 7
egg parasitoid, 1340
entomopathogenic nematode, 7
filth fly, 998
Fopius arisanus, 817
Heterorhabditis, 7
house fly, 593
insecticide, 1340
Muscidifurax raptorellus, 593, 998
Nasonia vitripennis, 593, 998
natural control, 617
Nosema pyrausta, 617
Ostrinia nubilalis, 617
parasitoid, 998
pesticides, 1075
poultry, 593, 998
predation, 617
Steinernema, 7
sublethal, 1075
toxicity, 1075
Trichogramma dendrolimi, 1340
virulence, 7
biological insecticides,
Acephate, 112
Bacillus thuringiensis variety *kurstaki*, 112
parasitism, 112
Rhyacionia frustrana, 112
tebufenozide, 112
black cutworm,
Agrotis ipsilon, 1183
endophyte, 1183
perennial ryegrass, 1183
Blattella germanica, 208, 499, 680, 686, 694, 1209, 1229
AChE, 499
attractants, 208, 686
bait, 1229
baits, 208, 686
chlorpyrifos, 499
cockroach, 686
coprophagy, 1229
cuticular penetration, 1209
esterase, 499
fipronil, 680
German cockroach, 1209
horizontal toxicant transfer, 680, 1229
permethrin, 694
pharmacokinetics, 1209
propoxur metabolism, 1209
resistance, 499, 694
Supella longipalpa, 686
synergists, 1209
translocation, 680
traps, 208

blueberry maggot,
Caylussacia spp., 1378
geographic distribution, 1378
host plants, 1378
Rhagoletis mendax, 1378
Vaccinium spp., 1378
Bollgard, 86, 1445
B.t. cotton, 86
Bollgard II, 1445
bollworm, 86
cotton, 1445
genetically modified, 1445
Helicoverpa zea, 86, 1445
insecticides, 86
Bollgard II,
Bollgard, 1445
cotton, 1445
genetically modified, 1445
Helicoverpa zea, 1445
bollworm,
B.t. cotton, 86
Bacillus thuringiensis, 76
Bollgard, 86
Helicoverpa zea, 76, 86
Heliothis virescens, 76
insecticide resistance, 76, 86
tobacco budworm, 76
Bombus impatiens, 462, 609
Apis mellifera, 609
bumble bee, 462, 609
greenhouse, 462
honey bee, 609
lowbush blueberry, 609
pollination, 609
pollination levels, 462
tomato, 462
Bombus terrestris,
imidacloprid, 623
seed dressing, 623
side effects, 623
sunflower, 623
Bonagota cranaodes,
Lepidoptera, 911
mating disruption, 911
monitoring trap, 911
sex pheromone, 911
Tortricidae, 911
Boophilus annulatus,
ivermectin, 1622
sustained release bolus, 1622
borate,
arsenic, 215
Incisitermes snyderi, 215
Spinosad, 215
borates,
Cryptotermes brevis, 915
drywood termite, 915
imidacloprid, 915
nuptial chamber, 915
silica gel, 915
breeding,
Apis mellifera, 1319
resistance, 1319
selection, 1319
Varroa destructor, 1319
bumble bee, 172, 609
Apis mellifera, 609
Bombus impatiens, 462
greenhouse, 172, 462
honey bee, 609
lowbush blueberry, 609
pollination, 172, 609
pollination levels, 462
tomato, 462
tomato quality, 172
Busseola fusca,
Cameroon, 1434
Eldana saccharina, 1434
maize stem borers, 1434
Mussidia nigrivenella, 1434
Sesamia calamistis, 1434

C

cabbage,
action threshold, 190
integrated pest management, 190
Trichoplusia ni, 190
cabbage maggot,
cabbage root fly, 942
crucifers, 942
Delia radicum, 942
host-plant resistance, 942
cabbage root fly,
cabbage maggot, 942
crucifers, 942
Delia radicum, 942
host-plant resistance, 942
cade oil,
alpha-ionol, 39
Bactrocera latifrons, 39
parapheromone, 39
caffeic acid,
Cylas formicarius elegantulus, 1285
nitrogen, 1285
plant resistance, 1285
resin glycoside, 1285
sweetpotato weevil, 1285
California,
distribution, 291
grapes, 291
mealybug, 291
Pseudococcus maritimus, 291
sampling, 291
Callosobruchus maculatus,
carbohydrate binding, 1280
cathepsin L, 1280
lectin, 1280
protease, 1280
protease inhibitor, 1280
Cameroon,
Busseola fusca, 1434
Eldana saccharina, 1434
maize stem borers, 1434
Mussidia nigrivenella, 1434
Sesamia calamistis, 1434
cantaloupe,
Bemisia argentifolii, 180
economic injury levels, 180
Carabidae,
ecotoxicology, 60
halofenozide, 60
Harpalus pennsylvanicus, 60
imidacloprid, 60
turfgrass, 60
carbaryl, 123, 929, 1428
areawide pest management, 929
consumption, 1428
cucurbitacin, 1428
Diabrotica virgifera virgifera, 929, 1428
insecticide resistance, 929
Ixodes scapularis, 123
mortality, 1428
nontarget effects, 123
carbohydrate binding,
Callosobruchus maculatus, 1280
cathepsin L, 1280
lectin, 1280
protease, 1280
protease inhibitor, 1280
carbon dioxide,
attractant, 860
Elasmopalpus lignosellus, 860
carboxylesterase,
cytochrome P450-dependent monooxygenase, 706
organophosphate insecticides, 706
Oryzaephilus surinamensis, 706
resistance, 706
catalase,
Diuraphis noxia, 743
peroxidase, 743
polyphenol oxidase, 743
protein content, 743
Rhopalosiphum padi, 743
cathepsin L,
Callosobruchus maculatus, 1280
carbohydrate binding, 1280
lectin, 1280
protease, 1280
protease inhibitor, 1280
cations,
enhancement, 1
Lymantria dispar, 1
viral activity, 1
cattle,
grazing cattle weight gains, 780
stable flies, 780
Stomoxys calcitrans, 780
CC trap,
Bemisia, 47

Bemisia argentifolii, 47
suction trap, 47
yellow sticky card trap, 47
cedrene,
Coptotermes formosanus, 1532
insect repellent, 1532
natural products, 1532
Vetiveria zizanioides, 1532
Ceratitis capitata, 817, 898, 971, 989, 1413
amplified fragment-length polymorphism, 989
areawide control, 817
attractants, 898
bait sprays, 817
biological control, 817
Ceratitis rosa, 989
DNA probes, 989
early detection, 971
food attractants, 971
Fopius arisanus, 817
ginger root oil, 1413
hosts, 971
Mediterranean fruit fly, 898
silver staining, 989
sterile insect technique, 1413
tephritids, 989
trapping, 971
Ceratitis rosa,
amplified fragment-length polymorphism, 989
Ceratitis capitata, 989
DNA probes, 989
silver staining, 989
tephritids, 989
cereal leaf beetle,
defoliation, 634
Oulema melanopu, 634
wheat, 634
yield, 634
Cerotoma trifurcata, 1459, 1594
bean leaf beetle, 1459
Diabrotica virgifera virgifera, 1594
glabrous, 1459
host plant resistance, 1594
Popillia japonica, 1594
pubescence, 1459
soybeans, 1594
trichome, 1459
chemical ecology,
Dendroctonus, 1113
Hylastes, 1113
Hylobius, 1113
Pinus, 1113
root insects, 1113
chemical insecticides,
apple, 68
Bacillus thuringiensis, 68
ecotoxicology, 68
population dynamics, 68
spiders, 68
Chilo suppressalis,
Bacillus thuringiensis, 271
cry1Ab gene, 271
resistance evaluation, 271
Scirpophaga incertulas, 271
transgenic rice, 271
China, 162, 445
elm leaf beetle, 162
Japanese beetle, 445
Popillia japonica, 445
preference, 445
Pyrrhalta luteola, 162
temperate, 445
temperate regions, 162
Ulmus, 162, 445
chitin synthesis inhibitors,
bait, 506
control, 506
subterranean termite, 506
chlorpyrifos, 486, 499, 524
acetylcholinesterase insensitivity, 524
AChE, 499
Blattella germanica, 499
esterase, 499
imidacloprid, 486
insecticide resistance, 524
methomyl, 524
nonrepellent termiticides, 486
resistance, 499
Reticulitermes flavipes, 486
Spodoptera exigua, 524
Choristoneura rosaceana, 468, 673
apple, 468
apples, 673
exposure, 468
insect growth regulator, 673
obliquebanded leafroller, 673
residue, 468
tebufenozide, 468, 673
Chortodes rufostrigata,
Claviceps purpurea, 1471
disease samples, 1471
grass seed, 1471
insect samples, 1471
Poa pratensis, 1471
circular statistics,
dispersal, 831
flight patterns, 831
Frankliniella occidentalis, 831
nectarine, 831
western flower thrips, 831
Citrullus lanatus,
Acalymma vittatum, 892
Diabrotica undecimpunctata, 892
Hawkesbury watermelon, 892
southern corn rootworm, 892
xanthene dye, 892
Citrus clementina,
Aphis gossypii, 439
economic injury level, 439
economic threshold, 439
Claviceps purpurea,
Chortodes rufostrigata, 1471
disease samples, 1471
grass seed, 1471
insect samples, 1471
Poa pratensis, 1471
Clavigralla tomentosicollis,
African yam bean, 724
insect bioassays, 724
lectins, 724
Sphenostylis stenocarpa, 724
cockroach,
attractants, 686
baits, 686
Blattella germanica, 686
Supella longipalpa, 686
codling moth,
apple, 1511
Cydia pomonella, 1511
overwintering, 1511
pear, 1511
pest management, 1511
Coleomegilla maculata,
biological control, 617
natural control, 617
Nosema pyrausta, 617
Ostrinia nubilalis, 617
predation, 617
Coleoptera, 367, 552, 844
Beauveria bassiana, 367
diatomaceous earth, 367
enemy free space, 844
fitness, 552
insecticide, 552
Ips pini, 844
malathion resistance, 552
mycopathogen, 367
phloem quality, 844
Pinus resinosa, 844
Scolytidae, 844
selection, 552
stored grain, 367
synergism, 367
Tribolium castaneum, 552
colony elimination,
automated termite monitoring system, 1518
datalogger, 1518
Isoptera, 1518
Colorado potato beetle, 204, 344, 572
Beauveria bassiana, 344
cyromazine, 344
host plant resistance, 572
imidacloprid, 344
Leptinotarsa decemlineata, 572
potato, 572
Solanum, 572
starvation, 344
synergism, 344
tomato, 204

trap, 204
compensation,
cotton, 388
pest management, 388
square removal, 388
composite index,
fitness, 826
fluctuating asymmetry, 826
mass rearing, 826
quality, 826
Trichogramma nr. *brassicae*, 826
compost,
bioassay, 1308
house fly, 1308
larval nutrition, 1308
Musca domestica, 1308
poultry manure, 1308
Conotrachelus nenuphar,
aggregation pheromone, 1386
fruit volatiles, 1386
monitoring traps, 1386
plum curculio, 1386
consumption,
carbaryl, 1428
cucurbitacin, 1428
Diabrotica virgifera virgifera, 1428
mortality, 1428
control,
bait, 506
chitin synthesis inhibitors, 506
subterranean termite, 506
controlled atmosphere,
Liposcelis bostrychophila, 1296
plant essential oil, 1296
toxicity, 1296
controlled atmospheres,
methyl bromide alternative, 1302
Plodia interpunctella, 1302
Rhyzopertha dominica, 1302
Tribolium castaneum, 1302
coprophagy,
bait, 1229
Blattella germanica, 1229
horizontal toxicant transfer, 1229
Coptotermes acinaciformis,
Coptotermes lacteus, 1602
Lincoln index, 1602
mark-recapture, 1602
Neutral Red, 1602
Nile Blue A, 1602
Coptotermes formosanus, 516, 1193, 1217, 1398, 1532
cedrene, 1532
deterrence, 516
feeding-preference, 516
foraging behavior, 1193
insect repellent, 1532
insecticide susceptibility, 1217
natural products, 1532
resistance, 1217
Reticulitermes flavipes, 1398
Reticulitermes virginicus, 1217
search pattern, 1193
subterranean termite, 516
subterranean termites, 1193
termite, 1217
termite behavior, 1398
termite foraging, 1398
termite tunneling, 1398
Veticeria zizanioides, 1532
wood, 516
Coptotermes lacteus,
Coptotermes acinaciformis, 1602
Lincoln index, 1602
mark-recapture, 1602
Neutral Red, 1602
Nile Blue A, 1602
corn, 98, 106
crop rotation, 98
Diabrotica virgifera virgifera, 98, 106
sampling, 98
soybeans, 98
threshold, 98
yield, 106
corn earworm,
Bacillus thuringiensis, 1268
sublethal effects, 1268
transgenic crops, 1268
corn rootworm,
adult mortality, 599
Diabrotica barberi, 599
Diabrotica virgifera virgifera, 599
dispersal, 599
larval survival, 599
Cosmopolites sordidus,
Azadirachta indica, 449
banana, 449
neem, 449
Cotinis nitida,
green June beetle, 628
organic fertilizer, 628
pastures, 628
white grubs, 628
cotton, 27, 388, 640, 1445
aphid–plant interactions, 27
Aphis gossypii, 27
Bollgard, 1445
Bollgard II, 1445
compensation, 388
deposit, 640
droplet size, 640
fecundity, 27
genetically modified, 1445
Helicoverpa zea, 1445
Heliothis virescens, 640
insecticide, 640
morphology, 27
nitrogen fertilization, 27
pest management, 388
square removal, 388
volumetric application rate, 640
cranberries,
(*E*)-11-tetradecenyl acetate, 658
mating disruption, 658
Sparganothis fruitworm, 658
Vaccinium macrocarpon, 658
crop damage,
Inopus rubriceps, 1097
sugarcane, 1097
crop rotation,
corn, 98
Diabrotica virgifera virgifera, 98
sampling, 98
soybeans, 98
threshold, 98
cross-resistance,
Drosophila melanogaster, 223
genetic variation, 223
insecticide resistance, 223
natural populations, 223
crucifers,
cabbage maggot, 942
cabbage root fly, 942
Delia radicum, 942
host-plant resistance, 942
cry1Ab gene,
Bacillus thuringiensis, 271
Chilo suppressalis, 271
resistance evaluation, 271
Scirpophaga incertulas, 271
transgenic rice, 271
Cry1Ac, 248, 752, 1237
B. thuringiensis cotton, 1237
Bacillus thuringiensis, 248, 1237
Cry2Ab, 752
inheritance, 248
Lepidoptera, 752
Pectinophora gossypiella, 248, 1237
resistance, 1237
resistance management, 248
toxicity, 752
transgenic *Bacillus thuringiensis* cotton, 752
transgenic cotton, 248
Cry1C,
Bacillus thuringiensis, 1547
Plutella xylostella, 1547
resistance, 1547
transgenic plants, 1547
Cry2Ab,
Cry1Ac, 752
Lepidoptera, 752
toxicity, 752
transgenic *Bacillus thuringiensis* cotton, 752

Cry9C,
Bacillus thuringiensis, 397
Diatrea grandiosella, 397
European corn borer, 397
Ostrinia nubilalis, 397
southwestern corn borer, 397
Cryptophlebia, 1005, 1292
alternative treatment efficacy, 1005
Bactrocera dorsalis, 1005
disinfestations, 1292
heat treatment, 1292
pest risk management, 1005
quarantine pest, 1292
quarantine treatment, 1005
Cryptophlebia illepida,
delayed mating, 785
demography, 785
life tables, 785
mating disruption, 785
Cryptotermes brevis,
borates, 915
drywood termite, 915
imidacloprid, 915
nuptial chamber, 915
silica gel, 915
cucumber,
esterase, 381
glutathione *S*-transferase, 381
maize, 381
pesticide susceptibility, 381
Tetranychus urticae, 381
cucurbitacin,
carbaryl, 1428
consumption, 1428
Diabrotica virgifera virgifera, 1428
mortality, 1428
Culex pipiens,
Bacillus sphaericus, 920
Bacillus thuringiensis, 920
microbial control, 920
population variation, 920
cultural control, 1485, 1526
aromatic cedar mulch, 1526
Delia antiqua, 1485
Delia radicum, 1485
landscaping, 1526
Linepithema humile, 1526
mechanical control, 1485
Tapinoma sessile, 1526
cultural practices,
host plant resistance, 963
Lissorhoptrus oryzophilus, 963
Oryza sativa, 963
resistance to infestation, 963
tolerance, 963
Curculio caryae,
biological control, 7
entomopathogenic nematode, 7
Heterorhabditis, 7
Steinernema, 7
virulence, 7
cuticular penetration,
Blattella germanica, 1209
German cockroach, 1209
pharmacokinetics, 1209
propoxur metabolism, 1209
synergists, 1209
Cydia pomonella, 264, 1511
apple, 1511
codling moth, 1511
insecticide bioassay, 264
insecticide resistance, 264
methoxyfenozide, 264
overwintering, 1511
pear, 1511
pest management, 1511
tebufenozide, 264
Cylas formicarius elegantulus,
caffeic acid, 1285
nitrogen, 1285
plant resistance, 1285
resin glycoside, 1285
sweetpotato weevil, 1285
cypermethrin, 868, 874
Noctuidae, 868, 874
permethrin, 868
pharmacokinetics, 874
S,S,S-tri-*n*-butyl phosphorotrithioate, 874
synergism, 868, 874
toxicity, 868
cyromazine,
Beauveria bassiana, 344
Colorado potato beetle, 344
imidacloprid, 344
starvation, 344
synergism, 344
cytochrome p450,
esterases, 253
Helicoverpa armigera, 253
India, 253
cytochrome P450,
inheritance, 1577
microfluorometric assay, 1577
mitochondrial electron transport inhibitor-acaricides, 1577
cytochrome p450,
nerve insensitivity, 253
pyrethroid resistance, 253
cytochrome P450,
resistance, 1577
Tetranychus urticae, 1577
cytochrome P450-dependent monooxygenase,
carboxylesterase, 706
organophosphate insecticides, 706
Oryzaephilus surinamensis, 706
resistance, 706

D

Dacus ciliatus,
detoxification, 1059
organophosphates, 1059
pesticides, 1059
pyrethroids, 1059
Tephritidae, 1059
Daktulosphaira vitifoliae,
grape, 138
season, 138
Vitis, 138
datalogger,
automated termite monitoring system, 1518
colony elimination, 1518
Isoptera, 1518
deet,
deet-insensitive mutant, 1584
Drosophila melanogaster, 1584
repellency, 1584
defoliation,
cereal leaf beetle, 634
Oulema melanopu, 634
wheat, 634
yield, 634
delayed mating,
Cryptophlebia illepida, 785
demography, 785
life tables, 785
mating disruption, 785
Delia antiqua,
cultural control, 1485
Delia radicum, 1485
mechanical control, 1485
Delia radicum, 942, 1485
cabbage maggot, 942
cabbage root fly, 942
crucifers, 942
cultural control, 1485
Delia antiqua, 1485
host-plant resistance, 942
mechanical control, 1485
demography,
Cryptophlebia illepida, 785
delayed mating, 785
life tables, 785
mating disruption, 785
Dendroctonus,
chemical ecology, 1113
Hylastes, 1113
Hylobius, 1113
Pinus, 1113
root insects, 1113
deployment strategies,
behavioral control, 905
imidacloprid-treated spheres, 905
Rhagoletis mendax, 905
deposit,
cotton, 640
droplet size, 640
Heliothis virescens, 640

insecticide, 640
volumetric application rate, 640
Dermanyssus gallinae,
mite, 1617
parasite, 1617
poultry, 1617
sample size, 1617
Taylor's power law, 1617
Dermestes maculatus,
Alphitobius diaperinus., 1610
hide beetle, 1610
lesser mealworm, 1610
Litter beetles, 1610
mechanical barrier, 1610
detection,
Diaprepes, 853
grubs, 853
Phyllophaga, 853
deterrence,
Coptotermes formosanus, 516
feeding-preference, 516
subterranean termite, 516
wood, 516
detoxification,
Dacus ciliatus, 1059
organophosphates, 1059
pesticides, 1059
pyrethroids, 1059
Tephritidae, 1059
Diabrotica barberi,
adult mortality, 599
corn rootworm, 599
Diabrotica virgifera virgifera, 599
dispersal, 599
larval survival, 599
Diabrotica undecimpunctata,
Acalymma vittatum, 892
Citrullus lanatus, 892
Hawkesbury watermelon, 892
southern corn rootworm, 892
xanthene dye, 892
Diabrotica virgifera virgifera, 98, 106, 599, 929, 1428, 1594
adult mortality, 599
areawide pest management, 929
carbaryl, 929, 1428
Cerotoma trifurcata, 1594
consumption, 1428
corn, 98, 106
corn rootworm, 599
crop rotation, 98
cucurbitacin, 1428
Diabrotica barberi, 599
dispersal, 599
host plant resistance, 1594
insecticide resistance, 929
larval survival, 599
mortality, 1428
Popillia japonica, 1594
sampling, 98
soybeans, 98, 1594
threshold, 98
yield, 106
Diabrotica virgifera virgifera,
host plant resistance, 529
population genetics, 529
resistance management, 529
simulation model, 529
Diachasmimorpha longicaudata,
Mediterranean fruit fly, 1419
Mexican fruit fly, 1419
phloxine B, 1419
phototoxic dye, 1419
West Indian fruit fly, 1419
Diadegma insulare, 14, 541
Microplitis plutellae, 14
parasitism, 14
permethrin, 541
Plutella xylostella, 14, 541
searching behavior, 14
diapause, 935, 1012
Bacillus thuringiensis, 935, 1012
fitness costs, 935
heat units, 1012
Pectinophora gossypiella, 935, 1012
resistance management, 1012
transgenic cotton, 935, 1012
Diaprepes,
detection, 853
grubs, 853
Phyllophaga, 853
diatomaceous earth,
Beauveria bassiana, 367
Coleoptera, 367
mycopathogen, 367
stored grain, 367
synergism, 367
Diatraea saccharalis, 1464, 1553
insecticide resistance, 1553
sugarcane, 1464, 1553
tebufenozide, 1464, 1553
Diatrea grandiosella,
Bacillus thuringiensis, 397
Cry9C, 397
European corn borer, 397
Ostrinia nubilalis, 397
southwestern corn borer, 397
diflubenzuron,
inheritance, 1243
Lucilia cuprina, 1243
monooxygenase, 1243
resistance, 1243
disease samples,
Chortodes rufostrigata, 1471
Claviceps purpurea, 1471
grass seed, 1471
insect samples, 1471
Poa pratensis, 1471
disinfestations,
Cryptophlebia, 1292
heat treatment, 1292
quarantine pest, 1292
dispersal, 315, 599, 698, 831
adult mortality, 599
Bacillus thuringiensis, 315, 698
circular statistics, 831
corn rootworm, 599
Diabrotica barberi, 599
Diabrotica virgifera virgifera, 599
flight patterns, 831
Frankliniella occidentalis, 831
geographical information system, 315
insecticide resistance, 698
larval survival, 599
local adaptation, 698
nectarine, 831
Pectinophora gossypiella, 315
resistance management, 315
transgenic cotton, 315
transgenic crops, 698
western flower thrips, 831
dispersion,
sequential sampling, 766
sugarcane, 766
sugarcane borer, 766
distribution,
California, 291
grapes, 291
mealybug, 291
Pseudococcus maritimus, 291
sampling, 291
Diuraphis noxia,
catalase, 743
peroxidase, 743
polyphenol oxidase, 743
protein content, 743
Rhopalosiphum padi, 743
DNA probes,
amplified fragment-length polymorphism, 989
Ceratitis capitata, 989
Ceratitis rosa, 989
silver staining, 989
tephritids, 989
droplet size,
cotton, 640
deposit, 640
Heliothis virescens, 640
insecticide, 640
volumetric application rate, 640
Drosophila melanogaster, 223, 1584
cross-resistance, 223
deet, 1584
deet-insensitive mutant, 1584
genetic variation, 223
insecticide resistance, 223
natural populations, 223
repellency, 1584

drywood termite,
borates, 915
Cryptotermes brevis, 915
imidacloprid, 915
nuptial chamber, 915
silica gel, 915
dual-toxin,
Bacilllus thuringiensis quantification, 1589
ELISA, 1589
transgenic cotton varieties, 1589

E

(*E*)-11-tetradecenyl acetate,
cranberries, 658
mating disruption, 658
Sparganothis fruitworm, 658
Vaccinium macrocarpon, 658
early detection,
Ceratitis capitata, 971
food attractants, 971
hosts, 971
trapping, 971
ecology,
Anthonomus grandis, 1405
herbivory, 1405
host plants, 1405
economic injury,
feeding behavior, 33
Malus domestica, 33
economic injury level, 439, 1162
Aphis gossypii, 439
Citrus clementina, 439
economic threshold, 439
Ostrinia nubilalis, 1162
potato, 1162
yield, 1162
economic injury levels,
Bemisia argentifolii, 180
cantaloupe, 180
economic threshold, 439, 1104
Aphis gossypii, 439
Citrus clementina, 439
economic injury level, 439
Helicoverpa armigera, 1104
plant scouting, 1104
tomatoes, 1104
ecotoxicology, 60, 68
apple, 68
Bacillus thuringiensis, 68
Carabidae, 60
chemical insecticides, 68
halofenozide, 60
Harpalus pennsylvanicus, 60
imidacloprid, 60
population dynamics, 68
spiders, 68
turfgrass, 60
efficacy,
flumethrin, 806
fluvalinate, 806
persistence, 806
resistance, 806
Varroa jacobsoni, 806
egg parasitoid,
biological control, 1340
insecticide, 1340
Trichogramma dendrolimi, 1340
Elasmopalpus lignosellus,
attractant, 860
carbon dioxide, 860
Eldana saccharina,
Busseola fusca, 1434
Cameroon, 1434
maize stem borers, 1434
Mussidia nigrivenella, 1434
Sesamia calamistis, 1434
ELISA,
Bacillus thuringiensis quantification, 1589
dual-toxin, 1589
transgenic cotton varieties, 1589
elm leaf beetle,
China, 162
Pyrrhalta luteola, 162
temperate regions, 162
Ulmus, 162
Empoasca fabae, 93, 950
alfalfa, 93, 950
glandular trichomes, 950
host plant resistance, 950
Medicago sativa, 950
phloem translocation, 93
potato leafhopper, 950
endophyte,
Agrotis ipsilon, 1183
black cutworm, 1183
perennial ryegrass, 1183
enemy free space,
Coleoptera, 844
Ips pini, 844
phloem quality, 844
Pinus resinosa, 844
Scolytidae, 844
enhancement,
cations, 1
Lymantria dispar, 1
viral activity, 1
entomopathogenic nematode,
biological control, 7
Curculio caryae, 7
Heterorhabditis, 7
Steinernema, 7
virulence, 7
enzyme-linked immunosorbent assay,
Bacillus thuringiensis quantification, 284
transgenic varieties, 284
essential oil extracts,
acaricide, 167
Artemisia absinthium, 167
Tanacetum vulgare, 167
twospotted spider mite, 167
esterase, 381, 499, 714
AChE, 499
Blattella germanica, 499
chlorpyrifos, 499
cucumber, 381
glutathione S-transferase, 381
Hematobia irritans irritans, 714
insecticide resistance, 714
maize, 381
pesticide susceptibility, 381
resistance, 499
Tetranychus urticae, 381
esterases, 253, 884
cytochrome p450, 253
Helicoverpa armigera, 253
India, 253
inhibition studies, 884
nerve insensitivity, 253
Noctuidae, 884
pyrethroid resistance, 253
S,S,S-tri-*n*-butyl phosphorotrithioate, 884
Euphorbia pulcherrima,
Bemisia argentifolii, 666
insecticide efficacy, 666
irrigation method, 666
European corn borer,
Bacillus thuringiensis, 397
Cry9C, 397
Diatrea grandiosella, 397
Ostrinia nubilalis, 397
southwestern corn borer, 397
Euschistus conspersus,
aggregation pheromone, 1500
Hemiptera, 1500
orchard, 1500
Pentatomidae, 1500
trapping, 1500
Euschistus servus,
Acrosternum hilare, 403
Nezara viridula, 403
exotic pest,
bark beetle, 422
Scolytidae, 422
Tomicus piniperda, 422
exposure, 468, 772
apple, 468
beetles, 772
Choristoneura rosaceana, 468
hydroprene, 772
morphological effects, 772
residue, 468
tebufenozide, 468
toxicity, 772
treated surfaces, 772

F

F_1 generation,
Heliothis virescens, 362

insectary, 362
rearing, 362
Serratia marcescens, 362
transmission, 362
favonoid,
a1-sh2, 564
Helicoverpa zea, 564
marker-assisted selection, 564
Zea mays, 564
fecundity, 27, 1344
aphid–plant interactions, 27
Aphis gossypii, 27
beneficial insect, 1344
cotton, 27
intrinsic rate of increase, 1344
morphology, 27
nitrogen fertilization, 27
Podisus maculiventris, 1344
predator, 1344
rearing, 1344
feeding behavior,
economic injury, 33
Malus domestica, 33
feeding-preference,
Coptotermes formosanus, 516
deterrence, 516
subterranean termite, 516
wood, 516
field activity,
*Ac*MNPV, 1037
*Af*MNPV, 1037
formulation, 1037
Lepidoptera, 1037
solar stability, 1037
filth fly,
biological control, 998
Muscidifurax raptorellus, 998
Nasonia vitripennis, 998
parasitoid, 998
poultry, 998
fipronil, 680, 811
Blattella germanica, 680
German yellowjacket, 811
horizontal toxicant transfer, 680
Patagonia, 811
pest control, 811
toxic baiting, 811
translocation, 680
Vespula germanica, 811
fitness, 552, 826
Coleoptera, 552
composite index, 826
fluctuating asymmetry, 826
insecticide, 552
malathion resistance, 552
mass rearing, 826
quality, 826
selection, 552
Tribolium castaneum, 552
Trichogramma nr. *brassicae*, 826
fitness costs,
Bacillus thuringiensis, 935
diapause, 935
Pectinophora gossypiella, 935
transgenic cotton, 935
flight patterns,
circular statistics, 831
dispersal, 831
Frankliniella occidentalis, 831
nectarine, 831
western flower thrips, 831
flight room,
Apis mellifera, 793
foraging, 793
insect rearing, 793
insect vision, 793
floriculture,
arthropods, 430
bedding plants, 430
greenhouse, 430
hypoxia, 430
fluctuating asymmetry,
composite index, 826
fitness, 826
mass rearing, 826
quality, 826
Trichogramma nr. *brassicae*, 826
flumethrin,
efficacy, 806
fluvalinate, 806
persistence, 806
resistance, 806
Varroa jacobsoni, 806
fluvalinate, 806, 1324
Apis mellifera, 1324
efficacy, 806
flumethrin, 806
honey bee, 1324
persistence, 806
resistance, 806
treatment threshold, 1324
Varroa destructor, 1324
Varroa jacobsoni, 806, 1324
food attractants,
Ceratitis capitata, 971
early detection, 971
hosts, 971
trapping, 971
Fopius arisanus,
areawide control, 817
bait sprays, 817
biological control, 817
Ceratitis capitata, 817
foraging,
Apis mellifera, 793
flight room, 793
insect rearing, 793
insect vision, 793
foraging behavior, 850, 1193
Coptotermes formosanus, 1193
Hymenoptera, 850
search pattern, 1193
social wasps, 850
subterranean termites, 1193
Vespula germanica, 850
forest,
ants, 761
bait traps, 761
Formicidae, 761
pitfall traps, 761
Formicidae,
ants, 761
bait traps, 761
forest, 761
pitfall traps, 761
formulation,
*Ac*MNPV, 1037
*Af*MNPV, 1037
field activity, 1037
Lepidoptera, 1037
solar stability, 1037
Frankliniella occidentalis, 579, 831, 984
circular statistics, 831
dispersal, 831
flight patterns, 831
greenhouse cucumber, 579
nectarine, 831
Platynota stultana, 984
Pseudococcus maritimus
Tetranychus urticae, 984
quarantine, 984
sequential sampling, 579
Tetranychus pacificus, 984
Vitis vinifera, 984
western flower thrips, 831
fruit volatiles,
aggregation pheromone, 1386
Conotrachelus nenuphar, 1386
monitoring traps, 1386
plum curculio, 1386
fumonisin,
Bacillus thuringiensis, 1067
Fusarium, 1067
Helicoverpa, 1067
Ostrinia, 1067
Fusarium,
Bacillus thuringiensis, 1067
fumonisin, 1067
Helicoverpa, 1067
Ostrinia, 1067

G

Gaylussacia spp.,
blueberry maggot, 1378
geographic distribution, 1378
host plants, 1378
Rhagoletis mendax, 1378
Vaccinium spp., 1378
generation mean analysis,
inbred lines, 277
maize, 277
plant resistance, 277

Sesamia nonagrioides, 277
genetic variation,
cross-resistance, 223
Drosophila melanogaster, 223
insecticide resistance, 223
natural populations, 223
genetically engineered crops,
Bacillus thuringiensis, 730
Ostrinia nubilalis, 730
Zea mays, 730
genetically modified,
Bollgard, 1445
Bollgard II, 1445
cotton, 1445
Helicoverpa zea, 1445
Geocoris punctipes,
bioassay, 55
insecticide, 55
Orius insidiosus, 55
geographic distribution,
blueberry maggot, 1378
Gaylussacia spp., 1378
host plants, 1378
Rhagoletis mendax, 1378
Vaccinium spp., 1378
geographical information system,
Bacillus thuringiensis, 315
dispersal, 315
Pectinophora gossypiella, 315
resistance management, 315
transgenic cotton, 315
geostatistics,
acoustic sampling, 1452
soil insects, 1452
German cockroach,
Blattella germanica, 1209
cuticular penetration, 1209
pharmacokinetics, 1209
propoxur metabolism, 1209
synergists, 1209
German yellowjacket,
fipronil, 811
Patagonia, 811
pest control, 811
toxic baiting, 811
Vespula germanica, 811
ginger root oil,
Ceratitis capitata, 1413
sterile insect technique, 1413
glabrous,
bean leaf beetle, 1459
Cerotoma trifurcata, 1459
pubescence, 1459
trichome, 1459
glandular trichomes,
alfalfa, 950
Empoasca fabae, 950
host plant resistance, 950
Medicago sativa, 950
potato leafhopper, 950
glufosinate-ammonium,
herbicide, 157
predatory insect, 157
predatory mite, 157
Tetranychus urticae, 157
toxicity, 157
glutathione *S*-transferase,
cucumber, 381
esterase, 381
maize, 381
pesticide susceptibility, 381
Tetranychus urticae, 381
glycoalkaloids,
host-plant resistance, 1260
leptines, 1260
Leptinotarsa decemlineata, 1260
potato, 1260
goatgrass,
antibiosis, 558
antixenosis, 558
Schizaphis graminum, 558
tolerance, 558
wheat, 558
grape,
Daktulosphaira vitifoliae, 138
season, 138
Vitis, 138
grapes,
California, 291
distribution, 291
mealybug, 291
Pseudococcus maritimus, 291
sampling, 291
Grapholita molesta, 233, 476
azinphosmethyl, 233
bioassay, 233
peach, 476
pest control, 476
resistance monitoring, 233
synergism, 233
grass seed,
Chortodes rufostrigata, 1471
Claviceps purpurea, 1471
disease samples, 1471
insect samples, 1471
Poa pratensis, 1471
grazing cattle weight gains,
cattle, 780
stable flies, 780
Stomoxys calcitrans, 780
green June beetle,
Cotinis nitida, 628
organic fertilizer, 628
pastures, 628
white grubs, 628
greenbug,
insecticide, 547
resistance, 547
sorghum, 547
wheat, 547
greenhouse, 172, 430, 462, 1492
Aphis gossypii, 1492
arthropods, 430
bedding plants, 430
Bombus impatiens, 462
bumble bee, 172, 462
floriculture, 430
hibiscus, 1492
horticulture, 1492
hypoxia, 430
pest management, 1492
pollination, 172
pollination levels, 462
tomato, 462
tomato quality, 172
wash, 1492
greenhouse cucumber,
Frankliniella occidentalis, 579
sequential sampling, 579
growth inhibition,
1,8-cineole, 979
antifeedant, 979
toxicity, 979
Tribolium castaneum, 979
grubs,
detection, 853
Diaprepes, 853
Phyllophaga, 853
gyrase inhibitor,
nutrition, 21
Pimpla turionellae, 21

H

halofenozide, 60, 145
ants, 145
Carabidae, 60
ecotoxicology, 60
Harpalus pennsylvanicus, 60
imidacloprid, 60, 145
Japanese beetles, 145
predators, 145
turfgrass, 60, 145
Harpalus pennsylvanicus,
Carabidae, 60
ecotoxicology, 60
halofenozide, 60
imidacloprid, 60
turfgrass, 60
Hawkesbury watermelon,
Acalymma vittatum, 892
Citrullus lanatus, 892
Diabrotica undecimpunctata, 892
southern corn rootworm, 892
xanthene dye, 892
heat treatment,
Cryptophlebia, 1292
disinfestations, 1292
quarantine pest, 1292
heat units,
Bacillus thuringiensis, 1012
diapause, 1012
Pectinophora gossypiella, 1012
resistance management, 1012
transgenic cotton, 1012

helicopter application, 1085, 1090
abamectin, 1090
avocado, 1085, 1090
sabadilla, 1085
Scirtothrips perseae, 1085, 1090
spray volumes, 1085, 1090
Helicoverpa,
Bacillus thuringiensis, 1067
fumonisin, 1067
Fusarium, 1067
Ostrinia, 1067
Helicoverpa armigera, 253, 1104
cytochrome p450, 253
economic threshold, 1104
esterases, 253
India, 253
nerve insensitivity, 253
plant scouting, 1104
pyrethroid resistance, 253
tomatoes, 1104
Helicoverpa zea, 76, 86, 197, 564, 1445
a1-sh2, 564
B.t. cotton, 86
Bacillus thuringiensis, 76, 197
Bollgard, 86, 1445
Bollgard II, 1445
bollworm, 76, 86
cotton, 1445
favonoid, 564
genetically modified, 1445
Heliothis virescens, 76
insecticide resistance, 76
insecticides, 86
marker-assisted selection, 564
Ostrinia nubilalis, 197
tobacco budworm, 76
transgenic sweet corn, 197
Zea mays, 564
Heliothis virescens, 76, 362, 640
Bacillus thuringiensis, 76
bollworm, 76
cotton, 640
deposit, 640
droplet size, 640
F_1 generation, 362
Helicoverpa zea, 76
insectary, 362
insecticide, 640
insecticide resistance, 76
rearing, 362
Serratia marcescens, 362
tobacco budworm, 76
transmission, 362
volumetric application rate, 640
Hematobia irritans irritans,
esterase, 714
insecticide resistance, 714
Hemiptera,
aggregation pheromone, 1500
Euschistus conspersus, 1500
orchard, 1500
Pentatomidae, 1500
trapping, 1500
herbicide,
glufosinate-ammonium, 157
predatory insect, 157
predatory mite, 157
Tetranychus urticae, 157
toxicity, 157
herbivory,
Anthonomus grandis, 1405
ecology, 1405
host plants, 1405
Heterorhabditis,
biological control, 7
Curculio caryae, 7
entomopathogenic nematode, 7
Steinernema, 7
virulence, 7
hibiscus,
Aphis gossypii, 1492
greenhouse, 1492
horticulture, 1492
pest management, 1492
wash, 1492
hide beetle,
Alphitobius diaperinus., 1610
Dermestes maculatus, 1610
lesser mealworm, 1610
litter beetles, 1610
mechanical barrier, 1610
Homalodisca coagulata,
Nerium oleander, 1031
oleander leaf scorch, 1031
vector, 1031
honey bee, 332, 609, 1022, 1324
Acarapis woodi, 332
Apis mellifera, 332, 609, 1022, 1324
bee breeding, 332
Bombus impatiens, 609
bumble bee, 609
fluvalinate, 1324
host-parasite, 1022
lowbush blueberry, 609
mite resistance, 332
parasitic mite, 1022
pollination, 609
tracheal mite, 332
treatment threshold, 1324
Varroa destructor, 1022, 1324
Varroa jacobsoni, 1022, 1324
horizontal toxicant transfer, 680, 1229
bait, 1229
Blattella germanica, 680, 1229
coprophagy, 1229
fipronil, 680
translocation, 680
horticulture,
Aphis gossypii, 1492
greenhouse, 1492
hibiscus, 1492
pest management, 1492
wash, 1492
host plant resistance, 529, 572, 950, 958, 963, 1249, 1594
alfalfa, 950
antibiosis, 958
Apion godmani, 958
beet armyworm, 1249
Cerotoma trifurcata, 1594
Colorado potato beetle, 572
cultural practices, 963
Diabrotica virgifera virgifera, 529, 1594
Empoasca fabae, 950
glandular trichomes, 950
hypersensitivity, 958
Leptinotarsa decemlineata, 572
Lissorhoptrus oryzophilus, 963
Lycopersicon hirsutum, 1249
Medicago sativa, 950
Oryza sativa, 963
Phaseolus vulgaris, 958
Popillia japonica, 1594
population genetics, 529
potato, 572
potato leafhopper, 950
resistance management, 529
resistance to infestation, 963
sesquiterpene, 1249
simulation model, 529
Solanum, 572
soybeans, 1594
tolerance, 963
tomato, 1249
tomato fruitworm, 1249
host plant-mediated effect,
Bacillus thuringiensis, 1135
Malacosoma disstria, 1135
quaking aspen, 1135
sugar maple, 1135
host plants, 1378, 1405
Anthonomus grandis, 1405
blueberry maggot, 1378
ecology, 1405
Gaylussacia spp., 1378
geographic distribution, 1378
herbivory, 1405
Rhagoletis mendax, 1378
Vaccinium spp., 1378
host records,
Lepidoptera, 1128
Tachinidae, 1128
host-parasite,
Apis mellifera, 1022
honey bee, 1022
parasitic mite, 1022
Varroa destructor, 1022
Varroa jacobsoni, 1022

host-plant resistance, 942, 1260
cabbage maggot, 942
cabbage root fly, 942
crucifers, 942
Delia radicum, 942
glycoalkaloids, 1260
leptines, 1260
Leptinotarsa decemlineata, 1260
potato, 1260
hosts,
Ceratitis capitata, 971
early detection, 971
food attractants, 971
trapping, 971
house fly, 593, 1308
bioassay, 1308
biological control, 593
compost, 1308
larval nutrition, 1308
Musca domestica, 1308
Muscidifurax raptorellus, 593
Nasonia vitripennis, 593
poultry, 593
poultry manure, 1308
hybrid economic injury level,
aesthetic threshold, 1187
azalea lace bug, 1187
Rhododendron spp., 1187
Stephanitis pyrioides, 1187
Hydrophilicity/lipophilicity balance ratios,
hydrophobic bonding, 1170
mating disruption, 1170
microsprayer, 1170
nonspecific membrane disruption, 1170
pheromones, 1170
hydrophobic bonding,
hydrophilicity/lipophilicity balance ratios, 1170
mating disruption, 1170
microsprayer, 1170
nonspecific membrane disruption, 1170
pheromones, 1170
hydroprene,
beetles, 772
exposure, 772
morphological effects, 772
toxicity, 772
treated surfaces, 772
hygienic behavior,
Apis mellifera, 326
Varroa destructor mite resistance, 326
Hylastes,
chemical ecology, 1113
Dendroctonus, 1113
Hylobius, 1113
Pinus, 1113
root insects, 1113
Hylobius,
chemical ecology, 1113
Dendroctonus, 1113
Hylastes, 1113
Pinus, 1113
root insects, 1113
Hymenoptera,
foraging behavior, 850
social wasps, 850
Vespula germanica, 850
hypersensitivity,
antibiosis, 958
Apion godmani, 958
host plant resistance, 958
Phaseolus vulgaris, 958
hypoxia,
arthropods, 430
bedding plants, 430
floriculture, 430
greenhouse, 430

I

imidacloprid, 60, 145, 344, 357, 486, 492, 623, 915
ants, 145
aversion, 492
Beauveria bassiana, 344, 357
Bemisia argentifolii, 357
Bombus terrestris, 623
borates, 915
Carabidae, 60
chlorpyrifos, 486
Colorado potato beetle, 344
Cryptotermes brevis, 915
cyromazine, 344
drywood termite, 915
ecotoxicology, 60
halofenozide, 60, 145
Harpalus pennsylvanicus, 60
insect pathogens, 357
Japanese beetles, 145
learning, 492
microbial control, 357
nonrepellent termiticides, 486
nuptial chamber, 915
predators, 145
Premise, 492
repellency, 492
Reticulitermes flavipes, 486
Reticulitermes virginicus, 492
seed dressing, 623
side effects, 623
silica gel, 915
starvation, 344
sunflower, 623
synergism, 344
turfgrass, 60, 145
whiteflies, 357
imidacloprid-treated spheres,
behavioral control, 905
deployment strategies, 905
Rhagoletis mendax, 905
inbred lines,
generation mean analysis, 277
maize, 277
plant resistance, 277
Sesamia nonagrioides, 277
Incisitermes snyderi,
arsenic, 215
borate, 215
Spinosad, 215
India,
cytochrome p450, 253
esterases, 253
Helicoverpa armigera, 253
nerve insensitivity, 253
pyrethroid resistance, 253
Indian meal moth,
Angoumois grain moth, 586
stored grain, 586
transgenic grain, 586
indoxacarb,
insecticide toxicology tests, 1122
parasitoids, 1122
pyrethroids, 1122
Rhyacionia frustrana, 1122
Spinosad, 1122
inheritance, 248, 1243, 1577
Bacillus thuringiensis, 248
Cry1Ac, 248
cytochrome P450, 1577
diflubenzuron, 1243
Lucilia cuprina, 1243
microfluorometric assay, 1577
mitochondrial electron transport inhibitor-acaricides, 1577
monooxygenase, 1243
Pectinophora gossypiella, 248
resistance, 1243, 1577
resistance management, 248
Tetranychus urticae, 1577
transgenic cotton, 248
inhibition studies,
esterases, 884
Noctuidae, 884
S,S,S-tri-*n*-butyl phosphorotrithioate, 884
Inopus rubriceps,
crop damage, 1097
sugarcane, 1097
insect bioassays,
African yam bean, 724
Clavigralla tomentosicollis, 724
lectins, 724
Sphenostylis stenocarpa, 724
insect dispersal,
Ostrinia nubilalis, 1369
resistance management, 1369
insect growth regulator,
apples, 673
Choristoneura rosaceana, 673
obliquebanded leafroller, 673

tebufenozide, 673
insect pathogen,
Bacillus thuringiensis, 1361
bioassay, 1361
larval development, 1361
Nosema pyrausta, 1361
Ostrinia nubilalis, 1361
insect pathogens,
Beauveria bassiana, 357
Bemisia argentifolii, 357
imidacloprid, 357
microbial control, 357
whiteflies, 357
insect rearing,
Apis mellifera, 793
flight room, 793
foraging, 793
insect vision, 793
insect repellent,
cedrene, 1532
Coptotermes formosanus, 1532
natural products, 1532
Vetiveria zizanioides, 1532
insect samples,
Chortodes rufostrigata, 1471
Claviceps purpurea, 1471
disease samples, 1471
grass seed, 1471
Poa pratensis, 1471
insect vision,
Apis mellifera, 793
flight room, 793
foraging, 793
insect rearing, 793
insectary,
F_1 generation, 362
Heliothis virescens, 362
rearing, 362
Serratia marcescens, 362
transmission, 362
insecticidal control,
action threshold, 1332
Myzus persicae, 1332
potato leafroll virus, 1332
vector, 1332
insecticide, 55, 150, 547, 552, 640, 1340
Anastrepha ludens, 150
bioassay, 55
biological control, 1340
Coleoptera, 552
cotton, 640
deposit, 640
droplet size, 640
egg parasitoid, 1340
fitness, 552
Geocoris punctipes, 55
greenbug, 547
Heliothis virescens, 640
malathion resistance, 552
Mexican fruit fly, 150
Orius insidiosus, 55
photo active dye, 150
resistance, 547
selection, 552
sorghum, 547
Tribolium castaneum, 552
Trichogramma dendrolimi, 1340
volumetric application rate, 640
wheat, 547
insecticide bioassay,
Cydia pomonella, 264
insecticide resistance, 264
methoxyfenozide, 264
tebufenozide, 264
insecticide efficacy,
Bemisia argentifolii, 666
Euphorbia pulcherrima, 666
irrigation method, 666
insecticide resistance, 76, 223, 264, 524, 698, 714, 929, 1553, 1558
acetylcholinesterase insensitivity, 524
Aleurothrixus floccosus, 1558
areawide pest management, 929
Bacillus thuringiensis, 76, 698
bioassay, 1558
bollworm, 76
carbaryl, 929
chlorpyrifos, 524
cross-resistance, 223
Cydia pomonella, 264
Diabrotica virgifera virgifera, 929
Diatraea saccharalis, 1553
dispersal, 698
Drosophila melanogaster, 223
esterase, 714
genetic variation, 223
Helicoverpa zea, 76
Heliothis virescens, 76
Hematobia irritans irritans, 714
insecticide bioassay, 264
local adaptation, 698
methomyl, 524
methoxyfenozide, 264
natural populations, 223
Spodoptera exigua, 524
sugarcane, 1553
tebufenozide, 264, 1553
tobacco budworm, 76
transgenic crops, 698
insecticide susceptibility,
Coptotermes formosanus, 1217
resistance, 1217
Reticulitermes virginicus, 1217
termite, 1217
insecticide toxicology tests,
indoxacarb, 1122
parasitoids, 1122
pyrethroids, 1122
Rhyacionia frustrana, 1122
Spinosad, 1122
insecticides, 86, 1353
B.t. cotton, 86
Bollgard, 86
bollworm, 86
Helicoverpa zea, 86
mortality, 1353
Orius insidiosus, 1353
Sorghum bicolor, 1353
Zea mays, 1353
insecticide-treated spheres,
Rhagoletis mendax, 1142
Rhagoletis pomonella, 1142
integrated pest management,
action threshold, 190
cabbage, 190
Trichoplusia ni, 190
intrinsic rate of increase,
beneficial insect, 1344
fecundity, 1344
Podisus maculiventris, 1344
predator, 1344
rearing, 1344
introduced crop,
apical damage, 648
kenaf, 648
Lygus pratensis, 648
Lygus rugulipennis, 648
introduced species,
Agromyzidae, 1177
invasive species, 1177
molecular identification, 1177
pea leafminer, 1177
polymerase chain reaction-restriction fragment-length polymorphism, 1177
invasive species,
Agromyzidae, 1177
introduced species, 1177
molecular identification, 1177
pea leafminer, 1177
polymerase chain reaction-restriction fragment-length polymorphism, 1177
Ips pini,
Coleoptera, 844
enemy free space, 844
phloem quality, 844
Pinus resinosa, 844
Scolytidae, 844
irrigation method,
Bemisia argentifolii, 666
Euphorbia pulcherrima, 666
insecticide efficacy, 666
Isoptera,
automated termite monitoring system, 1518
colony elimination, 1518

datalogger, 1518
ivermectin,
Boophilus annulatus, 1622
sustained release bolus, 1622
Ixodes scapularis,
carbaryl, 123
nontarget effects, 123

J

Japanese beetle,
China, 445
Popillia japonica, 445
preference, 445
temperate, 445
Ulmus, 445
Japanese beetles,
ants, 145
halofenozide, 145
imidacloprid, 145
predators, 145
turfgrass, 145

K

kenaf,
apical damage, 648
introduced crop, 648
Lygus pratensis, 648
Lygus rugulipennis, 648

L

landscaping,
aromatic cedar mulch, 1526
cultural control, 1526
Linepithema humile, 1526
Tapinoma sessile, 1526
larval development,
Bacillus thuringiensis, 1361
bioassay, 1361
insect pathogen, 1361
Nosema pyrausta, 1361
Ostrinia nubilalis, 1361
larval nutrition,
bioassay, 1308
compost, 1308
house fly, 1308
Musca domestica, 1308
poultry manure, 1308
larval survival,
adult mortality, 599
corn rootworm, 599
Diabrotica barberi, 599
Diabrotica virgifera virgifera, 599
dispersal, 599
LC_{50},
Bacillus thuringiensis, 1135
host plant-mediated effect, 1135
Malacosoma disstria, 1135
quaking aspen, 1135
sugar maple, 1135
leafhopper,
oleander leaf scorch, 1506
Pierce's disease, 1506
sharpshooter, 1506
vector, 1506
Xylella fastidiosa, 1506
leafrollers,
Argyrotaenia citrana, 373
mating disruption, 373
Pandemis pyrusana, 373
pheromones, 373
learning,
aversion, 492
imidacloprid, 492
Premise, 492
repellency, 492
Reticulitermes virginicus, 492
lectin,
Callosobruchus maculatus, 1280
carbohydrate binding, 1280
cathepsin L, 1280
protease, 1280
protease inhibitor, 1280
lectins,
African yam bean, 724
Clavigralla tomentosicollis, 724
insect bioassays, 724
Sphenostylis stenocarpa, 724
Lepidoptera, 752, 911, 1037, 1128, 1564
AcMNPV, 1037
AfMNPV, 1037
Bacillus thuringiensis, 1564
Bonagota cranaodes, 911
Cry1Ac, 752
Cry2Ab, 752
field activity, 1037
formulation, 1037
host records, 1128
mating disruption, 911
monitoring trap, 911
Ostrinia nubilalis, 1564
resistance, 1564
sex pheromone, 911
solar stability, 1037
Tachinidae, 1128
Tortricidae, 911
toxicity, 752
transgenic *Bacillus thuringiensis* cotton, 752
transgenic corn, 1564
leptines,
glycoalkaloids, 1260
host-plant resistance, 1260
Leptinotarsa decemlineata, 1260
potato, 1260
Leptinotarsa decemlineata, 572, 1260
Colorado potato beetle, 572
glycoalkaloids, 1260
host plant resistance, 572
host-plant resistance, 1260
leptines, 1260
potato, 572, 1260
Solanum, 572
lesser mealworm,
Alphitobius diaperinus., 1610
Dermestes maculatus, 1610
hide beetle, 1610
litter beetles, 1610
mechanical barrier, 1610
life tables,
Cryptophlebia illepida, 785
delayed mating, 785
demography, 785
mating disruption, 785
Lincoln index,
Coptotermes acinaciformis, 1602
Coptotermes lacteus, 1602
mark-recapture, 1602
Neutral Red, 1602
Nile Blue A, 1602
Linepithema humile, 511, 1526
ants, 511
Argentine ant, 511
aromatic cedar mulch, 1526
bait, 511
cultural control, 1526
gel, 511
landscaping, 1526
liquid, 511
Tapinoma sessile, 1526
Liposcelis bostrychophila,
controlled atmosphere, 1296
plant essential oil, 1296
toxicity, 1296
liquid,
ants, 511
Argentine ant, 511
bait, 511
gel, 511
Linepithema humile, 511
Lissorhoptrus oryzophilus,
cultural practices, 963
host plant resistance, 963
Oryza sativa, 963
resistance to infestation, 963
tolerance, 963
Litter beetles,
Alphitobius diaperinus., 1610
Dermestes maculatus, 1610
hide beetle, 1610
lesser mealworm, 1610
mechanical barrier, 1610
local adaptation,
Bacillus thuringiensis, 698
dispersal, 698
insecticide resistance, 698
transgenic crops, 698
lowbush blueberry,
Apis mellifera, 609
Bombus impatiens, 609
bumble bee, 609

honey bee, 609
pollination, 609
Lucilia cuprina,
diflubenzuron, 1243
inheritance, 1243
monooxygenase, 1243
resistance, 1243
Lycopersicon hirsutum,
beet armyworm, 1249
host plant resistance, 1249
sesquiterpene, 1249
tomato, 1249
tomato fruitworm, 1249
Lygus pratensis,
apical damage, 648
introduced crop, 648
kenaf, 648
Lygus rugulipennis, 648
Lygus rugulipennis,
apical damage, 648
introduced crop, 648
kenaf, 648
Lygus pratensis, 648
Lymantria dispar,
cations, 1
enhancement, 1
viral activity, 1

M

M2R,
Agrotis ipsilon, 1045
baculovirus, 1045
optical brightener, 1045
maize, 277, 381
cucumber, 381
esterase, 381
generation mean analysis, 277
glutathione S-transferase, 381
inbred lines, 277
pesticide susceptibility, 381
plant resistance, 277
Sesamia nonagrioides, 277
Tetranychus urticae, 381
maize stem borers,
Busseola fusca, 1434
Cameroon, 1434
Eldana saccharina, 1434
Mussidia nigrivenella, 1434
Sesamia calamistis, 1434
Malacosoma disstria,
Bacillus thuringiensis, 1135
host plant-mediated effect, 1135
LC_{50}, 1135
quaking aspen, 1135
sugar maple, 1135
malathion resistance,
Coleoptera, 552
fitness, 552
insecticide, 552
selection, 552
Tribolium castaneum, 552
Malus domestica,
economic injury, 33
feeding behavior, 33
marker-assisted selection,
a1-sh2, 564
favonoid, 564
Helicoverpa zea, 564
Zea mays, 564
mark-recapture,
Coptotermes acinaciformis, 1602
Coptotermes lacteus, 1602
Lincoln index, 1602
Neutral Red, 1602
Nile Blue A, 1602
mass rearing,
composite index, 826
fitness, 826
fluctuating asymmetry, 826
quality, 826
Trichogramma nr. *brassicae*, 826
maternal effects,
Bacillus thuringiensis, 1571
Pectinophora gossypiella, 1571
resistance costs, 1571
transgenic cotton, 1571
mating disruption, 373, 658, 785, 911, 1170
(*E*)-11-tetradecenyl acetate, 658
Argyrotaenia citrana, 373
Bonagota cranaodes, 911
cranberries, 658
Cryptophlebia illepida, 785
delayed mating, 785
demography, 785
hydrophilicity/lipophilicity balance ratios, 1170
hydrophobic bonding, 1170
leafrollers, 373
Lepidoptera, 911
life tables, 785
microsprayer, 1170
monitoring trap, 911
nonspecific membrane disruption, 1170
Pandemis pyrusana, 373
pheromones, 373, 1170
sex pheromone, 911
Sparganothis fruitworm, 658
Tortricidae, 911
Vaccinium macrocarpon, 658
mealybug,
California, 291
distribution, 291
grapes, 291
Pseudococcus maritimus, 291
sampling, 291
mechanical barrier,
Alphitobius diaperinus., 1610
Dermestes maculatus, 1610
hide beetle, 1610
lesser mealworm, 1610
litter beetles, 1610
mechanical control,
cultural control, 1485
Delia antiqua, 1485
Delia radicum, 1485
Medicago sativa,
alfalfa, 950
Empoasca fabae, 950
glandular trichomes, 950
host plant resistance, 950
potato leafhopper, 950
Mediterranean fruit fly, 898, 1419
attractants, 898
Ceratitis capitata, 898
Diachasmimorpha longicaudata, 1419
Mexican fruit fly, 1419
phloxine B, 1419
phototoxic dye, 1419
West Indian fruit fly, 1419
methomyl,
acetylcholinesterase insensitivity, 524
chlorpyrifos, 524
insecticide resistance, 524
Spodoptera exigua, 524
methoxyfenozide,
Cydia pomonella, 264
insecticide bioassay, 264
insecticide resistance, 264
tebufenozide, 264
methyl bromide alternative,
controlled atmospheres, 1302
Plodia interpunctella, 1302
Rhyzopertha dominica, 1302
Tribolium castaneum, 1302
Mexican fruit fly, 150, 1419
Anastrepha ludens, 150
Diachasmimorpha longicaudata, 1419
insecticide, 150
Mediterranean fruit fly, 1419
phloxine B, 1419
photo active dye, 150
phototoxic dye, 1419
West Indian fruit fly, 1419
microbial control, 357, 920
Bacillus sphaericus, 920
Bacillus thuringiensis, 920
Beauveria bassiana, 357
Bemisia argentifolii, 357
Culex pipiens, 920
imidacloprid, 357
insect pathogens, 357
population variation, 920
whiteflies, 357
microfluorometric assay,
cytochrome P450, 1577
inheritance, 1577
mitochondrial electron

transport inhibitor-acaricides, 1577
resistance, 1577
Tetranychus urticae, 1577
Microplitis plutellae,
Diadegma insulare, 14
parasitism, 14
Plutella xylostella, 14
searching behavior, 14
microsprayer,
hydrophilicity/lipophilicity balance ratios, 1170
hydrophobic bonding, 1170
mating disruption, 1170
nonspecific membrane disruption, 1170
pheromones, 1170
mite,
Dermanyssus gallinae, 1617
parasite, 1617
poultry, 1617
sample size, 1617
Taylor's power law, 1617
mite resistance,
Acarapis woodi, 332
Apis mellifera, 332
bee breeding, 332
honey bee, 332
tracheal mite, 332
mitochondrial electron transport inhibitor-acaricides,
cytochrome P450, 1577
inheritance, 1577
microfluorometric assay, 1577
resistance, 1577
Tetranychus urticae, 1577
molecular identification,
Agromyzidae, 1177
introduced species, 1177
invasive species, 1177
pea leafminer, 1177
polymerase chain reaction-restriction fragment-length polymorphism, 1177
monitoring,
pepper maggot, 455
Tephritidae, 455
Zonosemata electa, 455
monitoring trap,
Bonagota cranaodes, 911
Lepidoptera, 911
mating disruption, 911
sex pheromone, 911
Tortricidae, 911
monitoring traps,
aggregation pheromone, 1386
Conotrachelus nenuphar, 1386
fruit volatiles, 1386
plum curculio, 1386
monooxygenase,
diflubenzuron, 1243
inheritance, 1243
Lucilia cuprina, 1243
resistance, 1243
morphological effects,
beetles, 772
exposure, 772
hydroprene, 772
toxicity, 772
treated surfaces, 772
morphology,
aphid–plant interactions, 27
Aphis gossypii, 27
cotton, 27
fecundity, 27
nitrogen fertilization, 27
mortality, 1353, 1428
carbaryl, 1428
consumption, 1428
cucurbitacin, 1428
Diabrotica virgifera virgifera, 1428
insecticides, 1353
Orius insidiosus, 1353
Sorghum bicolor, 1353
Zea mays, 1353
Musa acuminata,
banana, 302
quarantine, 302
tephritid fruit flies, 302
trapping, 302
Musca domestica,
bioassay, 1308
compost, 1308
house fly, 1308
larval nutrition, 1308
poultry manure, 1308
Muscidifurax raptorellus, 593, 998
biological control, 593, 998
filth fly, 998
house fly, 593
Nasonia vitripennis, 593, 998
parasitoid, 998
poultry, 593, 998
Mussidia nigrivenella,
Busseola fusca, 1434
Cameroon, 1434
Eldana saccharina, 1434
maize stem borers, 1434
Sesamia calamistis, 1434
mycopathogen,
Beauveria bassiana, 367
Coleoptera, 367
diatomaceous earth, 367
stored grain, 367
synergism, 367
Myzus persicae,
action threshold, 1332
insecticidal control, 1332
potato leafroll virus, 1332
vector, 1332

N

Nasonia vitripennis, 593, 998
biological control, 593, 998
filth fly, 998
house fly, 593
Muscidifurax raptorellus, 593, 998
parasitoid, 998
poultry, 593, 998
natural control,
biological control, 617
Coleomegilla maculata, 617
Nosema pyrausta, 617
Ostrinia nubilalis, 617
predation, 617
natural populations,
cross-resistance, 223
Drosophila melanogaster, 223
genetic variation, 223
insecticide resistance, 223
natural products,
cedrene, 1532
Coptotermes formosanus, 1532
insect repellent, 1532
Vetiveria zizanioides, 1532
nectarine,
circular statistics, 831
dispersal, 831
flight patterns, 831
Frankliniella occidentalis, 831
western flower thrips, 831
neem,
Azadirachta indica, 449
banana, 449
Cosmopolites sordidus, 449
Nerium oleander,
Homalodisca coagulata, 1031
oleander leaf scorch, 1031
vector, 1031
nerve insensitivity,
cytochrome p450, 253
esterases, 253
Helicoverpa armigera, 253
India, 253
pyrethroid resistance, 253
Neutral Red,
Coptotermes acinaciformis, 1602
Coptotermes lacteus, 1602
Lincoln index, 1602
mark-recapture, 1602
Nile Blue A, 1602
Nezara viridula,
Acrosternum hilare, 403
Euschistus servus, 403
Nile Blue A,
Coptotermes acinaciformis, 1602
Coptotermes lacteus, 1602
Lincoln index, 1602
mark-recapture, 1602
Neutral Red, 1602

nitrogen,
caffeic acid, 1285
Cylas formicarius elegantulus, 1285
plant resistance, 1285
resin glycoside, 1285
sweetpotato weevil, 1285
nitrogen fertilization,
aphid–plant interactions, 27
Aphis gossypii, 27
cotton, 27
fecundity, 27
morphology, 27
Noctuidae, 868, 874, 884
cypermethrin, 868, 874
esterases, 884
inhibition studies, 884
permethrin, 868
pharmacokinetics, 874
S,S,S-tri-*n*-butyl phosphorotrithioate, 874, 884
synergism, 868, 874
toxicity, 868
nonchemical control,
plant extracts, 1200
repellents, 1200
subterranean termites, 1200
nonrepellent termiticides,
chlorpyrifos, 486
imidacloprid, 486
Reticulitermes flavipes, 486
nonspecific membrane disruption,
hydrophilicity/lipophilicity balance ratios, 1170
hydrophobic bonding, 1170
mating disruption, 1170
microsprayer, 1170
pheromones, 1170
nontarget effects,
carbaryl, 123
Ixodes scapularis, 123
Nosema pyrausta, 617, 1361
Bacillus thuringiensis, 1361
bioassay, 1361
biological control, 617
Coleomegilla maculata, 617
insect pathogen, 1361
larval development, 1361
natural control, 617
Ostrinia nubilalis, 617, 1361
predation, 617
nuptial chamber,
borates, 915
Cryptotermes brevis, 915
drywood termite, 915
imidacloprid, 915
silica gel, 915
nursery crops,
ornamentals, 1151
Popillia japonica, 1151
scarab, 1151
white grubs, 1151
nutrition,
gyrase inhibitor, 21
Pimpla turionellae, 21

O

obliquebanded leafroller,
apples, 673
Choristoneura rosaceana, 673
insect growth regulator, 673
tebufenozide, 673
oleander leaf scorch, 1031, 1506
Homalodisca coagulata, 1031
leafhopper, 1506
Nerium oleander, 1031
Pierce's disease, 1506
sharpshooter, 1506
vector, 1031, 1506
Xylella fastidiosa, 1506
optical brightener,
Agrotis ipsilon, 1045
baculovirus, 1045
M2R, 1045
optical brighteners,
Spodoptera exigua, 339
virus enhancement, 339
orchard,
aggregation pheromone, 1500
Euschistus conspersus, 1500
Hemiptera, 1500
Pentatomidae, 1500
trapping, 1500
organic fertilizer,
Cotinis nitida, 628
green June beetle, 628
pastures, 628
white grubs, 628
organophosphate insecticides,
carboxylesterase, 706
cytochrome P450-dependent monooxygenase, 706
Oryzaephilus surinamensis, 706
resistance, 706
organophosphates,
Dacus ciliatus, 1059
detoxification, 1059
pesticides, 1059
pyrethroids, 1059
Tephritidae, 1059
Orius insidiosus, 55, 1353
bioassay, 55
Geocoris punctipes, 55
insecticide, 55
insecticides, 1353
mortality, 1353
Sorghum bicolor, 1353
Zea mays, 1353
ornamentals,
nursery crops, 1151
Popillia japonica, 1151
scarab, 1151
white grubs, 1151
Oryza sativa,
cultural practices, 963
host plant resistance, 963
Lissorhoptrus oryzophilus, 963
resistance to infestation, 963
tolerance, 963
Oryzaephilus surinamensis,
carboxylesterase, 706
cytochrome P450-dependent monooxygenase, 706
organophosphate insecticides, 706
resistance, 706
Ostrinia,
Bacillus thuringiensis, 1067
fumonisin, 1067
Fusarium, 1067
Helicoverpa, 1067
Ostrinia nubilalis, 197, 397, 617, 730, 1162, 1361, 1369, 1564
Bacillus thuringiensis, 197, 397, 730, 1361, 1564
bioassay, 1361
biological control, 617
Coleomegilla maculata, 617
Cry9C, 397
Diatraea grandiosella, 397
economic injury level, 1162
European corn borer, 397
genetically engineered crops, 730
Helicoverpa zea, 197
insect dispersal, 1369
insect pathogen, 1361
larval development, 1361
Lepidoptera, 1564
natural control, 617
Nosema pyrausta, 617, 1361
potato, 1162
predation, 617
resistance, 1564
resistance management, 1369
southwestern corn borer, 397
transgenic corn, 1564
transgenic sweet corn, 197
yield, 1162
Zea mays, 730
Oulema melanopa,
cereal leaf beetle, 634
defoliation, 634
wheat, 634
yield, 634
overwintering,
apple, 1511
codling moth, 1511
Cydia pomonella, 1511
pear, 1511
pest management, 1511

P
Pandemis pyrusana,
Argyrotaenia citrana, 373
leafrollers, 373
mating disruption, 373
pheromones, 373
parapheromone,
alpha-ionol, 39
Bactrocera latifrons, 39
cade oil, 39
parasite,
Dermanyssus gallinae, 1617
mite, 1617
poultry, 1617
sample size, 1617
Taylor's power law, 1617
parasitic mite,
Apis mellifera, 1022
honey bee, 1022
host-parasite, 1022
Varroa destructor, 1022
Varroa jacobsoni, 1022
parasitism, 14, 112
Acephate, 112
Bacillus thuringiensis variety *kurstaki*, 112
biological insecticides, 112
Diadegma insulare, 14
Microplitis plutellae, 14
Plutella xylostella, 14
Rhyacionia frustrana, 112
searching behavior, 14
tebufenozide, 112
parasitoid,
biological control, 998
filth fly, 998
Muscidifurax raptorellus, 998
Nasonia vitripennis, 998
poultry, 998
parasitoids,
indoxacarb, 1122
insecticide toxicology tests, 1122
pyrethroids, 1122
Rhyacionia frustrana, 1122
spinosad, 1122
pastures,
Cotinis nitida, 628
green June beetle, 628
organic fertilizer, 628
white grubs, 628
Patagonia,
fipronil, 811
German yellowjacket, 811
pest control, 811
toxic baiting, 811
Vespula germanica, 811
pea leafminer,
Agromyzidae, 1177
introduced species, 1177
invasive species, 1177
molecular identification, 1177
polymerase chain reaction-restriction fragment-length polymorphism, 1177
peach,
Grapholita molesta, 476
pest control, 476
pear,
apple, 1511
codling moth, 1511
Cydia pomonella, 1511
overwintering, 1511
pest management, 1511
Pectinophora gossypiella, 248, 315, 935, 1012, 1237, 1571
B. thuringiensis cotton, 1237
Bacillus thuringiensis, 248, 315, 935, 1012, 1237, 1571
Cry1Ac, 248, 1237
diapause, 935, 1012
dispersal, 315
fitness costs, 935
geographical information system, 315
heat units, 1012
inheritance, 248
maternal effects, 1571
resistance, 1237
resistance costs, 1571
resistance management, 248, 315, 1012
transgenic cotton, 248, 315, 935, 1012, 1571
Pentatomidae,
aggregation pheromone, 1500
Euschistus conspersus, 1500
Hemiptera, 1500
orchard, 1500
trapping, 1500
pepper maggot,
monitoring, 455
Tephritidae, 455
Zonosemata electa, 455
perennial ryegrass,
Agrotis ipsilon, 1183
black cutworm, 1183
endophyte, 1183
permethrin, 541, 694, 868
Blattella germanica, 694
cypermethrin, 868
Diadegma insulare, 541
Noctuidae, 868
Plutella xylostella, 541
resistance, 694
synergism, 868
toxicity, 868
peroxidase,
catalase, 743
Diuraphis noxia, 743
polyphenol oxidase, 743
protein content, 743
Rhopalosiphum padi, 743
persistence,
efficacy, 806
flumethrin, 806
fluvalinate, 806
resistance, 806
Varroa jacobsoni, 806
pest control, 476, 811
fipronil, 811
German yellowjacket, 811
Grapholita molesta, 476
Patagonia, 811
peach, 476
toxic baiting, 811
Vespula germanica, 811
pest management, 388, 1492, 1511
Aphis gossypii, 1492
apple, 1511
codling moth, 1511
compensation, 388
cotton, 388
Cydia pomonella, 1511
greenhouse, 1492
hibiscus, 1492
horticulture, 1492
overwintering, 1511
pear, 1511
square removal, 388
wash, 1492
pest risk management,
alternative treatment efficacy, 1005
Bactrocera dorsalis, 1005
Cryptophlebia, 1005
quarantine treatment, 1005
pesticide susceptibility,
cucumber, 381
esterase, 381
glutathione *S*-transferase, 381
maize, 381
Tetranychus urticae, 381
pesticides, 1059, 1075
bioassay, 1075
biological control, 1075
Dacus ciliatus, 1059
detoxification, 1059
organophosphates, 1059
pyrethroids, 1059
sublethal, 1075
Tephritidae, 1059
toxicity, 1075
pharmacokinetics, 874, 1209
Blattella germanica, 1209
cuticular penetration, 1209
cypermethrin, 874
German cockroach, 1209
Noctuidae, 874
propoxur metabolism, 1209
S,S,S-tri-*n*-butyl phosphorotrithioate, 874
synergism, 874
synergists, 1209
Phaseolus vulgaris,

antibiosis, 958
Apion godmani, 958
host plant resistance, 958
hypersensitivity, 958
pheromones, 373, 1170
Argyrotaenia citrana, 373
hydrophilicity/lipophilicity balance ratios, 1170
hydrophobic bonding, 1170
leafrollers, 373
mating disruption, 373, 1170
microsprayer, 1170
nonspecific membrane disruption, 1170
Pandemis pyrusana, 373
phloem quality,
Coleoptera, 844
enemy free space, 844
Ips pini, 844
Pinus resinosa, 844
Scolytidae, 844
phloem translocation,
alfalfa, 93
Empoasca fabae, 93
phloxine B,
Diachasmimorpha longicaudata, 1419
Mediterranean fruit fly, 1419
Mexican fruit fly, 1419
phototoxic dye, 1419
West Indian fruit fly, 1419
photo active dye,
Anastrepha ludens, 150
insecticide, 150
Mexican fruit fly, 150
phototoxic dye,
Diachasmimorpha longicaudata, 1419
Mediterranean fruit fly, 1419
Mexican fruit fly, 1419
phloxine B, 1419
West Indian fruit fly, 1419
Phyllophaga,
detection, 853
Diaprepes, 853
grubs, 853
Pierce's disease,
leafhopper, 1506
oleander leaf scorch, 1506
sharpshooter, 1506
vector, 1506
Xylella fastidiosa, 1506
Pimpla turionellae,
gyrase inhibitor, 21
nutrition, 21
Pinus,
chemical ecology, 1113
Dendroctonus, 1113
Hylastes, 1113
Hylobius, 1113
root insects, 1113
Pinus resinosa,
Coleoptera, 844
enemy free space, 844
Ips pini, 844
phloem quality, 844
Scolytidae, 844
pitfall traps,
ants, 761
bait traps, 761
forest, 761
Formicidae, 761
plant essential oil,
controlled atmosphere, 1296
Liposcelis bostrychophila, 1296
toxicity, 1296
plant extracts,
nonchemical control, 1200
repellents, 1200
subterranean termites, 1200
plant resistance, 277, 1285
caffeic acid, 1285
Cylas formicarius elegantulus, 1285
generation mean analysis, 277
inbred lines, 277
maize, 277
nitrogen, 1285
resin glycoside, 1285
Sesamia nonagrioides, 277
sweetpotato weevil, 1285
plant scouting,
economic threshold, 1104
Helicoverpa armigera, 1104
tomatoes, 1104
Platynota stultana,
Frankliniella occidentalis, 984
Pseudococcus maritimus
Tetranychus urticae, 984
quarantine, 984
Tetranychus pacificus, 984
Vitis vinifera, 984
Plodia interpunctella,
controlled atmospheres, 1302
methyl bromide alternative, 1302
Rhyzopertha dominica, 1302
Tribolium castaneum, 1302
plum curculio,
aggregation pheromone, 1386
Conotrachelus nenuphar, 1386
fruit volatiles, 1386
monitoring traps, 1386
Plutella xylostella, 14, 240, 541, 1547
Bacillus thuringiensis, 240, 1547
Cry1C, 1547
Diadegma insulare, 14, 541
Microplitis plutellae, 14
parasitism, 14
permethrin, 541
resistance, 240, 1547
searching behavior, 14
transgenic plants, 240, 1547
Poa pratensis,
Chortodes rufostrigata, 1471
Claviceps purpurea, 1471
disease samples, 1471
grass seed, 1471
insect samples, 1471
Podisus maculiventris,
beneficial insect, 1344
fecundity, 1344
intrinsic rate of increase, 1344
predator, 1344
rearing, 1344
pollination, 172, 609
Apis mellifera, 609
Bombus impatiens, 609
bumble bee, 172, 609
greenhouse, 172
honey bee, 609
lowbush blueberry, 609
tomato quality, 172
pollination levels,
Bombus impatiens, 462
bumble bee, 462
greenhouse, 462
tomato, 462
polymerase chain reaction-restriction fragment-length polymorphism,
Agromyzidae, 1177
introduced species, 1177
invasive species, 1177
molecular identification, 1177
pea leafminer, 1177
polyphenol oxidase,
catalase, 743
Diuraphis noxia, 743
peroxidase, 743
protein content, 743
Rhopalosiphum padi, 743
Popillia japonica, 445, 1151, 1594
Cerotoma trifurcata, 1594
China, 445
Diabrotica virgifera virgifera, 1594
host plant resistance, 1594
Japanese beetle, 445
nursery crops, 1151
ornamentals, 1151
preference, 445
scarab, 1151
soybeans, 1594
temperate, 445
Ulmus, 445
white grubs, 1151
population dynamics,
apple, 68
Bacillus thuringiensis, 68
chemical insecticides, 68
ecotoxicology, 68
spiders, 68
population genetics,

Diabrotica virgifera virgifera, 529
host plant resistance, 529
resistance management, 529
simulation model, 529
population variation,
Bacillus sphaericus, 920
Bacillus thuringiensis, 920
Culex pipiens, 920
microbial control, 920
potato, 572, 1162, 1260
Colorado potato beetle, 572
economic injury level, 1162
glycoalkaloids, 1260
host plant resistance, 572
host-plant resistance, 1260
leptines, 1260
Leptinotarsa decemlineata, 572, 1260
Ostrinia nubilalis, 1162
Solanum, 572
yield, 1162
potato leafhopper,
alfalfa, 950
Empoasca fabae, 950
glandular trichomes, 950
host plant resistance, 950
Medicago sativa, 950
potato leafroll virus,
action threshold, 1332
insecticidal control, 1332
Myzus persicae, 1332
vector, 1332
poultry, 593, 998, 1617
biological control, 593, 998
Dermanyssus gallinae, 1617
filth fly, 998
house fly, 593
mite, 1617
Muscidifurax raptorellus, 593, 998
Nasonia vitripennis, 593, 998
parasite, 1617
parasitoid, 998
sample size, 1617
Taylor's power law, 1617
poultry manure,
bioassay, 1308
compost, 1308
house fly, 1308
larval nutrition, 1308
Musca domestica, 1308
predation,
biological control, 617
Coleomegilla maculata, 617
natural control, 617
Nosema pyrausta, 617
Ostrinia nubilalis, 617
predator,
beneficial insect, 1344
fecundity, 1344
intrinsic rate of increase, 1344
Podisus maculiventris, 1344
rearing, 1344
predators,
ants, 145
halofenozide, 145
imidacloprid, 145
Japanese beetles, 145
turfgrass, 145
predatory insect,
glufosinate-ammonium, 157
herbicide, 157
predatory mite, 157
Tetranychus urticae, 157
toxicity, 157
predatory mite,
glufosinate-ammonium, 157
herbicide, 157
predatory insect, 157
Tetranychus urticae, 157
toxicity, 157
preference,
China, 445
Japanese beetle, 445
Popillia japonica, 445
temperate, 445
Ulmus, 445
Premise,
aversion, 492
imidacloprid, 492
learning, 492
repellency, 492
Reticulitermes virginicus, 492
propoxur metabolism,
Blattella germanica, 1209
cuticular penetration, 1209
German cockroach, 1209
pharmacokinetics, 1209
synergists, 1209
protease,
Callosobruchus maculatus, 1280
carbohydrate binding, 1280
cathepsin L, 1280
lectin, 1280
protease inhibitor, 1280
protease inhibitor,
Callosobruchus maculatus, 1280
carbohydrate binding, 1280
cathepsin L, 1280
lectin, 1280
protease, 1280
protein content,
catalase, 743
Diuraphis noxia, 743
peroxidase, 743
polyphenol oxidase, 743
Rhopalosiphum padi, 743
Pseudococcus maritimus,
California, 291
distribution, 291
grapes, 291
mealybug, 291
sampling, 291
Pseudococcus maritimus
Tetranychus urticae,
Frankliniella occidentalis, 984
Platynota stultana, 984
quarantine, 984
Tetranychus pacificus, 984
Vitis vinifera, 984
Pseudoplusia includens,
Bacillus thuringiensis, 1053
soybean looper, 1053
transgenic cotton, 1053
pubescence,
bean leaf beetle, 1459
Cerotoma trifurcata, 1459
glabrous, 1459
trichome, 1459
pyrethroid resistance,
cytochrome p450, 253
esterases, 253
Helicoverpa armigera, 253
India, 253
nerve insensitivity, 253
pyrethroids, 1059, 1122
Dacus ciliatus, 1059
detoxification, 1059
indoxacarb, 1122
insecticide toxicology tests, 1122
organophosphates, 1059
parasitoids, 1122
pesticides, 1059
Rhyacionia frustrana, 1122
Spinosad, 1122
Tephritidae, 1059
Pyrrhalta luteola,
China, 162
elm leaf beetle, 162
temperate regions, 162
Ulmus, 162

Q

quaking aspen,
Bacillus thuringiensis, 1135
host plant-mediated effect, 1135
LC_{50}, 1135
Malacosoma disstria, 1135
sugar maple, 1135
quarantine, 302, 984
banana, 302
Frankliniella occidentalis, 984
Musa acuminata, 302
Platynota stultana, 984
Pseudococcus maritimus
Tetranychus urticae, 984
tephritid fruit flies, 302
Tetranychus pacificus, 984
trapping, 302
Vitis vinifera, 984
quarantine pest,
Cryptophlebia, 1292
disinfestations, 1292

heat treatment, 1292
quarantine treatment,
alternative treatment efficacy, 1005
Bactrocera dorsalis, 1005
Cryptophlebia, 1005
pest risk management, 1005

R

rearing, 362, 1344
beneficial insect, 1344
F_1 generation, 362
fecundity, 1344
Heliothis virescens, 362
insectary, 362
intrinsic rate of increase, 1344
Podisus maculiventris, 1344
predator, 1344
Serratia marcescens, 362
transmission, 362
reflective mulch,
resistant cultivars, 1477
strawberry, 1477
tarnished plant bug, 1477
repellency, 492, 1584
aversion, 492
deet, 1584
deet-insensitive mutant, 1584
Drosophila melanogaster, 1584
imidacloprid, 492
learning, 492
Premise, 492
Reticulitermes virginicus, 492
repellents,
nonchemical control, 1200
plant extracts, 1200
subterranean termites, 1200
residue,
apple, 468
Choristoneura rosaceana, 468
exposure, 468
tebufenozide, 468
resin glycoside,
caffeic acid, 1285
Cylas formicarius elegantulus, 1285
nitrogen, 1285
plant resistance, 1285
sweetpotato weevil, 1285
resistance, 240, 499, 547, 694, 706, 806, 1217, 1237, 1243, 1319, 1547, 1564, 1577
AChE, 499
Apis mellifera, 1319
B. thuringiensis cotton, 1237
Bacillus thuringiensis, 240, 1237, 1547, 1564
Blattella germanica, 499, 694
breeding, 1319
carboxylesterase, 706
chlorpyrifos, 499
Coptotermes formosanus, 1217
Cry1Ac, 1237
Cry1C, 1547
cytochrome P450, 1577
cytochrome P450-dependent monooxygenase, 706
diflubenzuron, 1243
efficacy, 806
esterase, 499
flumethrin, 806
fluvalinate, 806
greenbug, 547
inheritance, 1243, 1577
insecticide, 547
insecticide susceptibility, 1217
Lepidoptera, 1564
Lucilia cuprina, 1243
microfluorometric assay, 1577
mitochondrial electron transport inhibitor-acaricides, 1577
monooxygenase, 1243
organophosphate insecticides, 706
Oryzaephilus surinamensis, 706
Ostrinia nubilalis, 1564
Pectinophora gossypiella, 1237
permethrin, 694
persistence, 806
Plutella xylostella, 240, 1547
Reticulitermes virginicus, 1217
selection, 1319
sorghum, 547
termite, 1217
Tetranychus urticae, 1577
transgenic corn, 1564
transgenic plants, 240, 1547
Varroa destructor, 1319
Varroa jacobsoni, 806
wheat, 547
resistance costs,
Bacillus thuringiensis, 1571
maternal effects, 1571
Pectinophora gossypiella, 1571
transgenic cotton, 1571
resistance evaluation,
Bacillus thuringiensis, 271
Chilo suppressalis, 271
cry1Ab gene, 271
Scirpophaga incertulas, 271
transgenic rice, 271
resistance management, 248, 315, 529, 1012, 1369
Bacillus thuringiensis, 248, 315, 1012
Cry1Ac, 248
Diabrotica virgifera virgifera, 529
diapause, 1012
dispersal, 315
geographical information system, 315
heat units, 1012
host plant resistance, 529
inheritance, 248
insect dispersal, 1369
Ostrinia nubilalis, 1369
Pectinophora gossypiella, 248, 315, 1012
population genetics, 529
simulation model, 529
transgenic cotton, 248, 315, 1012
resistance monitoring, 233, 1538
azinphosmethyl, 233
baseline susceptibility, 1538
Bemisia argentifolii, 1538
bioassay, 233
Grapholita molesta, 233
synergism, 233
whiteflies, 1538
resistance to infestation,
cultural practices, 963
host plant resistance, 963
Lissorhoptrus oryzophilus, 963
Oryza sativa, 963
tolerance, 963
resistant cultivars,
reflective mulch, 1477
strawberry, 1477
tarnished plant bug, 1477
Reticulitermes flavipes, 486, 1398
chlorpyrifos, 486
Coptotermes formosanus, 1398
imidacloprid, 486
nonrepellent termiticides, 486
termite behavior, 1398
termite foraging, 1398
termite tunneling, 1398
Reticulitermes virginicus, 492, 1217
aversion, 492
Coptotermes formosanus, 1217
imidacloprid, 492
insecticide susceptibility, 1217
learning, 492
Premise, 492
repellency, 492
resistance, 1217
termite, 1217
Rhagoletis mendax, 905, 1142, 1378
behavioral control, 905
blueberry maggot, 1378
deployment strategies, 905
Gaylussacia spp., 1378
geographic distribution, 1378
host plants, 1378
imidacloprid-treated spheres, 905
insecticide-treated spheres, 1142
Rhagoletis pomonella, 1142
Vaccinium spp., 1378
Rhagoletis pomonella,

insecticide-treated spheres, 1142
Rhagoletis mendax, 1142
Rhododendron spp., 129, 1187
aesthetic threshold, 1187
azalea lace bug, 129, 1187
hybrid economic injury level, 1187
Stephanitis pyrioides, 129, 1187
Rhopalosiphum padi, 410, 743
barley yellow dwarf, 410
catalase, 743
Diuraphis noxia, 743
peroxidase, 743
polyphenol oxidase, 743
protein content, 743
Rhopalosiphum rufiabdominalis, 410
Schizaphis graminum, 410
Sitobion avenae, 410
wheat, 410
Rhopalosiphum rufiabdominalis,
barley yellow dwarf, 410
Rhopalosiphum padi, 410
Schizaphis graminum, 410
Sitobion avenae, 410
wheat, 410
Rhyacionia frustrana, 112, 1122
Acephate, 112
Bacillus thuringiensis variety *kurstaki*, 112
biological insecticides, 112
indoxacarb, 1122
insecticide toxicology tests, 1122
parasitism, 112
parasitoids, 1122
pyrethroids, 1122
spinosad, 1122
tebufenozide, 112
Rhyzopertha dominica,
controlled atmospheres, 1302
methyl bromide alternative, 1302
Plodia interpunctella, 1302
Tribolium castaneum, 1302
root insects,
chemical ecology, 1113
Dendroctonus, 1113
Hylastes, 1113
Hylobius, 1113
Pinus, 1113

S

S,S,S-tri-*n*-butyl phosphorotrithioate, 874, 884
cypermethrin, 874
esterases, 884
inhibition studies, 884
Noctuidae, 874, 884
pharmacokinetics, 874
synergism, 874
sabadilla,
avocado, 1085
helicopter application, 1085
Scirtothrips perseae, 1085
spray volumes, 1085
sample size,
Dermanyssus gallinae, 1617
mite, 1617
parasite, 1617
poultry, 1617
Taylor's power law, 1617
sampling, 98, 291
California, 291
corn, 98
crop rotation, 98
Diabrotica virgifera virgifera, 98
distribution, 291
grapes, 291
mealybug, 291
Pseudococcus maritimus, 291
soybeans, 98
threshold, 98
scarab,
nursery crops, 1151
ornamentals, 1151
Popillia japonica, 1151
white grubs, 1151
Schizaphis graminum, 410, 558
antibiosis, 558
antixenosis, 558
barley yellow dwarf, 410
goatgrass, 558
Rhopalosiphum padi, 410
Rhopalosiphum rufiabdominalis, 410
Sitobion avenae, 410
tolerance, 558
wheat, 410, 558
Scirpophaga incertulas,
Bacillus thuringiensis, 271
Chilo suppressalis, 271
cry1Ab gene, 271
resistance evaluation, 271
transgenic rice, 271
Scirtothrips perseae, 1085, 1090
abamectin, 1090
avocado, 1085, 1090
helicopter application, 1085, 1090
sabadilla, 1085
spray volumes, 1085, 1090
Scolytidae, 422, 844
bark beetle, 422
Coleoptera, 844
enemy free space, 844
exotic pest, 422
Ips pini, 844
phloem quality, 844
Pinus resinosa, 844
Tomicus piniperda, 422
search pattern,
Coptotermes formosanus, 1193
foraging behavior, 1193
subterranean termites, 1193
searching behavior,
Diadegma insulare, 14
Microplitis plutellae, 14
parasitism, 14
Plutella xylostella, 14
season,
Daktulosphaira vitifoliae, 138
grape, 138
Vitis, 138
seed dressing,
Bombus terrestris, 623
imidacloprid, 623
side effects, 623
sunflower, 623
selection, 552, 1319
Apis mellifera, 1319
breeding, 1319
Coleoptera, 552
fitness, 552
insecticide, 552
malathion resistance, 552
resistance, 1319
Tribolium castaneum, 552
Varroa destructor, 1319
sequential sampling, 579, 766
dispersion, 766
Frankliniella occidentalis, 579
greenhouse cucumber, 579
sugarcane, 766
Sugarcane borer, 766
Serratia marcescens,
F_1 generation, 362
Heliothis virescens, 362
insectary, 362
rearing, 362
transmission, 362
Sesamia calamistis,
Busseola fusca, 1434
Cameroon, 1434
Eldana saccharina, 1434
maize stem borers, 1434
Mussidia nigrivenella, 1434
Sesamia nonagrioides,
generation mean analysis, 277
inbred lines, 277
maize, 277
plant resistance, 277
sesquiterpene,
beet armyworm, 1249
host plant resistance, 1249
Lycopersicon hirsutum, 1249
tomato, 1249
tomato fruitworm, 1249
sex pheromone,
Bonagota cranaodes, 911
Lepidoptera, 911
mating disruption, 911
monitoring trap, 911

Tortricidae, 911
sharpshooter,
leafhopper, 1506
oleander leaf scorch, 1506
Pierce's disease, 1506
vector, 1506
Xylella fastidiosa, 1506
side effects,
Bombus terrestris, 623
imidacloprid, 623
seed dressing, 623
sunflower, 623
silica gel,
borates, 915
Cryptotermes brevis, 915
drywood termite, 915
imidacloprid, 915
nuptial chamber, 915
silver staining,
amplified fragment-length polymorphism, 989
Ceratitis capitata, 989
Ceratitis rosa, 989
DNA probes, 989
tephritids, 989
simulation model,
Diabrotica virgifera virgifera, 529
host plant resistance, 529
population genetics, 529
resistance management, 529
Sitobion avenae,
barley yellow dwarf, 410
Rhopalosiphum padi, 410
Rhopalosiphum rufiabdominalis, 410
Schizaphis graminum, 410
wheat, 410
social wasps,
foraging behavior, 850
Hymenoptera, 850
Vespula germanica, 850
soil insects,
acoustic sampling, 1452
geostatistics, 1452
Solanum,
Colorado potato beetle, 572
host plant resistance, 572
Leptinotarsa decemlineata, 572
potato, 572
solar stability,
AcMNPV, 1037
AfMNPV, 1037
field activity, 1037
formulation, 1037
Lepidoptera, 1037
sorghum,
greenbug, 547
insecticide, 547
resistance, 547
wheat, 547
Sorghum bicolor,
insecticides, 1353
mortality, 1353
Orius insidiosus, 1353
Zea mays, 1353
southern corn rootworm,
Acalymma vittatum, 892
Citrullus lanatus, 892
Diabrotica undecimpunctata, 892
Hawkesbury watermelon, 892
xanthene dye, 892
southwestern corn borer,
Bacillus thuringiensis, 397
Cry9C, 397
Diatrea grandiosella, 397
European corn borer, 397
Ostrinia nubilalis, 397
soybean looper,
Bacillus thuringiensis, 1053
Pseudoplusia includens, 1053
transgenic cotton, 1053
soybeans, 98, 1594
Cerotoma trifurcata, 1594
corn, 98
crop rotation, 98
Diabrotica virgifera virgifera, 98, 1594
host plant resistance, 1594
Popillia japonica, 1594
sampling, 98
threshold, 98
Sparganothis fruitworm,
(*E*)-11-tetradecenyl acetate, 658
cranberries, 658
mating disruption, 658
Vaccinium macrocarpon, 658
Sphenostylis stenocarpa,
African yam bean, 724
Clavigralla tomentosicollis, 724
insect bioassays, 724
lectins, 724
spiders,
apple, 68
Bacillus thuringiensis, 68
chemical insecticides, 68
ecotoxicology, 68
population dynamics, 68
Spinosad, 215, 1122
arsenic, 215
borate, 215
Incisitermes snyderi, 215
indoxacarb, 1122
insecticide toxicology tests, 1122
parasitoids, 1122
pyrethroids, 1122
Rhyacionia frustrana, 1122
Spodoptera exigua, 339, 524
acetylcholinesterase insensitivity, 524
chlorpyrifos, 524
insecticide resistance, 524
methomyl, 524
optical brighteners, 339
virus enhancement, 339
spray volumes, 1085, 1090
abamectin, 1090
avocado, 1085, 1090
helicopter application, 1085, 1090
sabadilla, 1085
Scirtothrips perseae, 1085, 1090
square removal,
compensation, 388
cotton, 388
pest management, 388
stable flies,
cattle, 780
grazing cattle weight gains, 780
Stomoxys calcitrans, 780
starvation,
Beauveria bassiana, 344
Colorado potato beetle, 344
cyromazine, 344
imidacloprid, 344
synergism, 344
Steinernema,
biological control, 7
Curculio caryae, 7
entomopathogenic nematode, 7
Heterorhabditis, 7
virulence, 7
Stephanitis pyrioides, 129, 1187
aesthetic threshold, 1187
azalea lace bug, 129, 1187
hybrid economic injury level, 1187
Rhododendron spp., 129, 1187
sterile insect technique,
Ceratitis capitata, 1413
ginger root oil, 1413
Stomoxys calcitrans,
cattle, 780
grazing cattle weight gains, 780
stable flies, 780
stored grain, 367, 586
Angoumois grain moth, 586
Beauveria bassiana, 367
Coleoptera, 367
diatomaceous earth, 367
Indian meal moth, 586
mycopathogen, 367
synergism, 367
transgenic grain, 586
strawberry,
reflective mulch, 1477
resistant cultivars, 1477
tarnished plant bug, 1477
sublethal,
bioassay, 1075

biological control, 1075
pesticides, 1075
toxicity, 1075
sublethal effects,
Bacillus thuringiensis, 1268
corn earworm, 1268
transgenic crops, 1268
subterranean termite, 506, 516
bait, 506
chitin synthesis inhibitors, 506
control, 506
Coptotermes formosanus, 516
deterrence, 516
feeding-preference, 516
wood, 516
subterranean termites, 1193, 1200
Coptotermes formosanus, 1193
foraging behavior, 1193
nonchemical control, 1200
plant extracts, 1200
repellents, 1200
search pattern, 1193
suction trap,
Bemisia, 47
Bemisia argentifolii, 47
CC trap, 47
yellow sticky card trap, 47
sugar maple,
Bacillus thuringiensis, 1135
host plant-mediated effect, 1135
LC_{50}, 1135
Malacosoma disstria, 1135
quaking aspen, 1135
sugarcane, 766, 1097, 1464, 1553
crop damage, 1097
Diatraea saccharalis, 1464, 1553
dispersion, 766
Inopus rubriceps, 1097
insecticide resistance, 1553
sequential sampling, 766
sugarcane borer, 766
tebufenozide, 1464, 1553
sugarcane borer,
dispersion, 766
sequential sampling, 766
sugarcane, 766
sunflower,
Bombus terrestris, 623
imidacloprid, 623
seed dressing, 623
side effects, 623
Supella longipalpa,
attractants, 686
baits, 686
Blattella germanica, 686
cockroach, 686
sustained release bolus,
Boophilus annulatus, 1622
ivermectin, 1622
sweetpotato weevil,
caffeic acid, 1285
Cylas formicarius elegantulus, 1285
nitrogen, 1285
plant resistance, 1285
resin glycoside, 1285
synergism, 233, 344, 367, 868, 874
azinphosmethyl, 233
Beauveria bassiana, 344, 367
bioassay, 233
Coleoptera, 367
Colorado potato beetle, 344
cypermethrin, 868, 874
cyromazine, 344
diatomaceous earth, 367
Grapholita molesta, 233
imidacloprid, 344
mycopathogen, 367
Noctuidae, 868, 874
permethrin, 868
pharmacokinetics, 874
resistance monitoring, 233
S,S,S-tri-*n*-butyl phosphorotrithioate, 874
starvation, 344
stored grain, 367
toxicity, 868
synergists,
Blattella germanica, 1209
cuticular penetration, 1209
German cockroach, 1209
pharmacokinetics, 1209
propoxur metabolism, 1209

T

Tachinidae,
host records, 1128
Lepidoptera, 1128
Tanacetum vulgare,
acaricide, 167
Artemisia absinthium, 167
essential oil extracts, 167
twospotted spider mite, 167
Tapinoma sessile,
aromatic cedar mulch, 1526
cultural control, 1526
landscaping, 1526
Linepithema humile, 1526
tarnished plant bug,
reflective mulch, 1477
resistant cultivars, 1477
strawberry, 1477
Taylor's power law,
Dermanyssus gallinae, 1617
mite, 1617
parasite, 1617
poultry, 1617
sample size, 1617
tebufenozide, 112, 264, 468, 673, 1464, 1553
Acephate, 112
apple, 468
apples, 673
Bacillus thuringiensis variety *kurstaki*, 112
biological insecticides, 112
Choristoneura rosaceana, 468, 673
Cydia pomonella, 264
Diatraea saccharalis, 1464, 1553
exposure, 468
insect growth regulator, 673
insecticide bioassay, 264
insecticide resistance, 264, 1553
methoxyfenozide, 264
obliquebanded leafroller, 673
parasitism, 112
residue, 468
Rhyacionia frustrana, 112
sugarcane, 1464, 1553
temperate,
China, 445
Japanese beetle, 445
Popillia japonica, 445
preference, 445
Ulmus, 445
temperate regions,
China, 162
elm leaf beetle, 162
Pyrrhalta luteola, 162
Ulmus, 162
tephritid fruit flies,
banana, 302
Musa acuminata, 302
quarantine, 302
trapping, 302
Tephritidae, 455, 1059
Dacus ciliatus, 1059
detoxification, 1059
monitoring, 455
organophosphates, 1059
pepper maggot, 455
pesticides, 1059
pyrethroids, 1059
Zonosemata electa, 455
tephritids,
amplified fragment-length polymorphism, 989
Ceratitis capitata, 989
Ceratitis rosa, 989
DNA probes, 989
silver staining, 989
termite,
Coptotermes formosanus, 1217
insecticide susceptibility, 1217
resistance, 1217
Reticulitermes virginicus, 1217
termite behavior,
Coptotermes formosanus, 1398
Reticulitermes flavipes, 1398
termite foraging, 1398
termite tunneling, 1398

termite foraging,
Coptotermes formosanus, 1398
Reticulitermes flavipes, 1398
termite behavior, 1398
termite tunneling, 1398
termite tunneling,
Coptotermes formosanus, 1398
Reticulitermes flavipes, 1398
termite behavior, 1398
termite foraging, 1398
Tetranychus pacificus,
Frankliniella occidentalis, 984
Platynota stultana, 984
Pseudococcus maritimus *Tetranychus urticae*, 984
quarantine, 984
Vitis vinifera, 984
Tetranychus urticae, 157, 381, 1577
cucumber, 381
cytochrome P450, 1577
esterase, 381
glufosinate-ammonium, 157
glutathione *S*-transferase, 381
herbicide, 157
inheritance, 1577
maize, 381
microfluorometric assay, 1577
mitochondrial electron transport inhibitor-acaricides, 1577
pesticide susceptibility, 381
predatory insect, 157
predatory mite, 157
resistance, 1577
toxicity, 157
threshold,
corn, 98
crop rotation, 98
Diabrotica virgifera virgifera, 98
sampling, 98
soybeans, 98
tobacco budworm,
Bacillus thuringiensis, 76
bollworm, 76
Helicoverpa zea, 76
Heliothis virescens, 76
insecticide resistance, 76
tolerance, 358, 963
antibiosis, 358
antixenosis, 358
cultural practices, 963
goatgrass, 358
host plant resistance, 963
Lissorhoptrus oryzophilus, 963
Oryza sativa, 963
resistance to infestation, 963
Schizaphis graminum, 358
wheat, 358
tomato, 204, 462, 1249
beet armyworm, 1249
Bombus impatiens, 462
bumble bee, 462
Colorado potato beetle, 204
greenhouse, 462
host plant resistance, 1249
Lycopersicon hirsutum, 1249
pollination levels, 462
sesquiterpene, 1249
tomato fruitworm, 1249
trap, 204
tomato fruitworm,
beet armyworm, 1249
host plant resistance, 1249
Lycopersicon hirsutum, 1249
sesquiterpene, 1249
tomato, 1249
tomato quality,
bumble bee, 172
greenhouse, 172
pollination, 172
tomatoes,
economic threshold, 1104
Helicoverpa armigera, 1104
plant scouting, 1104
Tomicus piniperda,
bark beetle, 422
exotic pest, 422
Scolytidae, 422
Tortricidae,
Bonagota cranaodes, 911
Lepidoptera, 911
mating disruption, 911
monitoring trap, 911
sex pheromone, 911
toxic baiting,
fipronil, 811
German yellowjacket, 811
Patagonia, 811
pest control, 811
Vespula germanica, 811
toxicity, 157, 752, 772, 868, 979, 1075, 1296
1,8-cineole, 979
antifeedant, 979
beetles, 772
bioassay, 1075
biological control, 1075
controlled atmosphere, 1296
Cry1Ac, 752
Cry2Ab, 752
cypermethrin, 868
exposure, 772
glufosinate-ammonium, 157
growth inhibition, 979
herbicide, 157
hydroprene, 772
Lepidoptera, 752
Liposcelis bostrychophila, 1296
morphological effects, 772
Noctuidae, 868
permethrin, 868
pesticides, 1075
plant essential oil, 1296
predatory insect, 157
predatory mite, 157
sublethal, 1075
synergism, 868
Tetranychus urticae, 157
transgenic *Bacillus thuringiensis* cotton, 752
treated surfaces, 772
Tribolium castaneum, 979
tracheal mite,
Acarapis woodi, 332
Apis mellifera, 332
bee breeding, 332
honey bee, 332
mite resistance, 332
transgenic *Bacillus thuringiensis* cotton,
Cry1Ac, 752
Cry2Ab, 752
Lepidoptera, 752
toxicity, 752
transgenic corn,
Bacillus thuringiensis, 1564
Lepidoptera, 1564
Ostrinia nubilalis, 1564
resistance, 1564
transgenic cotton, 248, 315, 935, 1012, 1053, 1571
Bacillus thuringiensis, 248, 315, 935, 1012, 1053, 1571
Cry1Ac, 248
diapause, 935, 1012
dispersal, 315
fitness costs, 935
geographical information system, 315
heat units, 1012
inheritance, 248
maternal effects, 1571
Pectinophora gossypiella, 248, 315, 935, 1012, 1571
Pseudoplusia includens, 1053
resistance costs, 1571
resistance management, 248, 315, 1012
soybean looper, 1053
transgenic cotton varieties,
Bacillus thuringiensis quantification, 1589
dual-toxin, 1589
ELISA, 1589
transgenic crops, 698, 1268
Bacillus thuringiensis, 698, 1268
corn earworm, 1268
dispersal, 698
insecticide resistance, 698
local adaptation, 698
sublethal effects, 1268
transgenic grain,
Angoumois grain moth, 586
Indian meal moth, 586
stored grain, 586

transgenic plants, 240, 1547
Bacillus thuringiensis, 240, 1547
Cry1C, 1547
Plutella xylostella, 240, 1547
resistance, 240, 1547
transgenic rice,
Bacillus thuringiensis, 271
Chilo suppressalis, 271
cry1Ab gene, 271
resistance evaluation, 271
Scirpophaga incertulas, 271
transgenic sweet corn,
Bacillus thuringiensis, 197
Helicoverpa zea, 197
Ostrinia nubilalis, 197
transgenic varieties,
Bacillus thuringiensis quantification, 284
enzyme-linked immunosorbent assay, 284
translocation,
Blattella germanica, 680
fipronil, 680
horizontal toxicant transfer, 680
transmission,
F_1 generation, 362
Heliothis virescens, 362
insectary, 362
rearing, 362
Serratia marcescens, 362
trap,
Colorado potato beetle, 204
tomato, 204
trapping, 302, 971, 1500
aggregation pheromone, 1500
banana, 302
Ceratitis capitata, 971
early detection, 971
Euschistus conspersus, 1500
food attractants, 971
Hemiptera, 1500
hosts, 971
Musa acuminata, 302
orchard, 1500
Pentatomidae, 1500
quarantine, 302
tephritid fruit flies, 302
traps,
attractants, 208
baits, 208
Blattella germanica, 208
treated surfaces,
beetles, 772
exposure, 772
hydroprene, 772
morphological effects, 772
toxicity, 772
treatment threshold,
Apis mellifera, 1324
fluvalinate, 1324
honey bee, 1324
Varroa destructor, 1324
Varroa jacobsoni, 1324
Tribolium castaneum, 552, 979, 1302
1,8-cineole, 979
antifeedant, 979
Coleoptera, 552
controlled atmospheres, 1302
fitness, 552
growth inhibition, 979
insecticide, 552
malathion resistance, 552
methyl bromide alternative, 1302
Plodia interpunctella, 1302
Rhyzopertha dominica, 1302
selection, 552
toxicity, 979
Trichogramma dendrolimi,
biological control, 1340
egg parasitoid, 1340
insecticide, 1340
Trichogramma nr. *brassicae*,
composite index, 826
fitness, 826
fluctuating asymmetry, 826
mass rearing, 826
quality, 826
trichome,
bean leaf beetle, 1459
Cerotoma trifurcata, 1459
glabrous, 1459
pubescence, 1459
Trichoplusia ni,
action threshold, 190
cabbage, 190
integrated pest management, 190
turfgrass, 60, 145
ants, 145
Carabidae, 60
ecotoxicology, 60
halofenozide, 60, 145
Harpalus pennsylvanicus, 60
imidacloprid, 60, 145
Japanese beetles, 145
predators, 145
twospotted spider mite,
acaricide, 167
Artemisia absinthium, 167
essential oil extracts, 167
Tanacetum vulgare, 167

U

Ulmus, 162, 445
China, 162, 445
elm leaf beetle, 162
Japanese beetle, 445
Popillia japonica, 445
preference, 445
Pyrrhalta luteola, 162
temperate, 445
temperate regions, 162
urban forestry,
Anoplophora glabripennis, 116
urban forests, 116
urban forests,
Anoplophora glabripennis, 116
urban forestry, 116

V

Vaccinium macrocarpon,
(*E*)-11-tetradecenyl acetate, 658
cranberries, 658
mating disruption, 658
Sparganothis fruitworm, 658
Vaccinium spp.,
blueberry maggot, 1378
Gaylussacia spp., 1378
geographic distribution, 1378
host plants, 1378
Rhagoletis mendax, 1378
Varroa destructor, 1022, 1319, 1324
Apis mellifera, 1022, 1324
Apis mellifera, 1319
breeding, 1319
fluvalinate, 1324
honey bee, 1022, 1324
host-parasite, 1022
parasitic mite, 1022
resistance, 1319
selection, 1319
treatment threshold, 1324
Varroa jacobsoni, 1022, 1324
Varroa destructor mite resistance,
Apis mellifera, 326
hygienic behavior, 326
Varroa jacobsoni, 806, 1022, 1324
Apis mellifera, 1022, 1324
efficacy, 806
flumethrin, 806
fluvalinate, 806, 1324
honey bee, 1022, 1324
host-parasite, 1022
parasitic mite, 1022
persistence, 806
resistance, 806
treatment threshold, 1324
Varroa destructor, 1022, 1324
vector, 1031, 1332, 1506
action threshold, 1332
Homalodisca coagulata, 1031
insecticidal control, 1332
leafhopper, 1506
Myzus persicae, 1332
Nerium oleander, 1031
oleander leaf scorch, 1031, 1506
Pierce's disease, 1506
potato leafroll virus, 1332
sharpshooter, 1506

Xylella fastidiosa, 1506
Vespula germanica, 811, 850
fipronil, 811
foraging behavior, 850
German yellowjacket, 811
Hymenoptera, 850
Patagonia, 811
pest control, 811
social wasps, 850
toxic baiting, 811
Vetiveria zizanioides,
cedrene, 1532
Coptotermes formosanus, 1532
insect repellent, 1532
natural products, 1532
viral activity,
cations, 1
enhancement, 1
Lymantria dispar, 1
virulence,
biological control, 7
Curculio caryae, 7
entomopathogenic nematode, 7
Heterorhabditis, 7
Steinernema, 7
virus enhancement,
optical brighteners, 339
Spodoptera exigua, 339
Vitis,
Daktulosphaira vitifoliae, 138
grape, 138
season, 138
Vitis vinifera,
Frankliniella occidentalis, 984
Platynota stultana, 984
Pseudococcus maritimus
Tetranychus urticae, 984
quarantine, 984
Tetranychus pacificus, 984
volumetric application rate,
cotton, 640
deposit, 640
droplet size, 640
Heliothis virescens, 640
insecticide, 640

W

wash,
Aphis gossypii, 1492
greenhouse, 1492
hibiscus, 1492
horticulture, 1492
pest management, 1492
West Indian fruit fly,
Diachasmimorpha longicaudata, 1419
Mediterranean fruit fly, 1419
Mexican fruit fly, 1419
phloxine B, 1419
phototoxic dye, 1419
western flower thrips,
circular statistics, 831
dispersal, 831
flight patterns, 831
Frankliniella occidentalis, 831
nectarine, 831
wheat, 410, 547, 558, 634
antibiosis, 558
antixenosis, 558
barley yellow dwarf, 410
cereal leaf beetle, 634
defoliation, 634
goatgrass, 558
greenbug, 547
insecticide, 547
Oulema melanopu, 634
resistance, 547
Rhopalosiphum padi, 410
Rhopalosiphum rufiabdominalis, 410
Schizaphis graminum, 410, 558
Sitobion avenae, 410
sorghum, 547
tolerance, 558
yield, 634
white grubs, 628, 1151
Cotinis nitida, 628
green June beetle, 628
nursery crops, 1151
organic fertilizer, 628
ornamentals, 1151
pastures, 628
Popillia japonica, 1151
scarab, 1151
whiteflies, 357, 1538
baseline susceptibility, 1538
Beauveria bassiana, 357
Bemisia argentifolii, 357, 1538
imidacloprid, 357
insect pathogens, 357
microbial control, 357
resistance monitoring, 1538
wood,
Coptotermes formosanus, 516
deterrence, 516
feeding-preference, 516
subterranean termite, 516

X

xanthene dye,
Acalymma vittatum, 892
Citrullus lanatus, 892
Diabrotica undecimpunctata, 892
Hawkesbury watermelon, 892
southern corn rootworm, 892
Xylella fastidiosa,
leafhopper, 1506
oleander leaf scorch, 1506
Pierce's disease, 1506
sharpshooter, 1506
vector, 1506

Y

yellow sticky card trap,
Bemisia, 47
Bemisia argentifolii, 47
CC trap, 47
suction trap, 47
yield, 106, 634, 1162
cereal leaf beetle, 634
corn, 106
defoliation, 634
Diabrotica virgifera virgifera, 106
economic injury level, 1162
Ostrinia nubilalis, 1162
Oulema melanopu, 634
potato, 1162
wheat, 634

Z

Zea mays, 564, 730, 1353
a1-sh2, 564
Bacillus thuringiensis, 730
favonoid, 564
genetically engineered crops, 730
Helicoverpa zea, 564
insecticides, 1353
marker-assisted selection, 564
mortality, 1353
Orius insidiosus, 1353
Ostrinia nubilalis, 730
Sorghum bicolor, 1353
Zonosemata electa,
monitoring, 455
pepper maggot, 455
Tephritidae, 455